

DETERMINATION OF REAL BOTT MANIFOLDS

ADMI NAZRA

ABSTRACT. A real Bott manifold is obtained as the orbit space of the n -torus T^n by the free action of an elementary abelian 2-group $(\mathbb{Z}_2)^n$. This paper deals with the classification of 5-dimensional real Bott manifolds and study certain type of n -dimensional real Bott manifolds ($n \geq 6$).

INTRODUCTION

A real Bott tower is described as a sequence of $\mathbb{RP}^1$ -bundles of height n which is the real restriction to a Bott tower introduced in [1]. The total space of such a sequence is called a *real Bott manifold*. From the viewpoint of group actions, an n -dimensional real Bott manifold is the quotient of the n -dimensional torus $T^n = S^1 \times \cdots \times S^1$ by the product $(\mathbb{Z}_2)^n$ of cyclic group of order 2. A *Bott matrix* A of size n is an upper triangular matrix whose diagonal entries are 1 and the other entries are either 1 or 0. By the definition, the number of distinct Bott matrices of size n is $2^{\frac{1}{2}(n^2-n)}$. The free action of $(\mathbb{Z}_2)^n$ on T^n can be expressed by each row of the Bott matrix A whose orbit space $M(A) = T^n/(\mathbb{Z}_2)^n$ is the real Bott manifold. It is easy to see that $M(A)$ is a compact euclidean space form (Riemannian flat manifold). Then we can apply the Bieberbach theorem [5] to classify real Bott manifolds. Using this theorem, the classification of real Bott manifolds up to dimension 4 has been obtained in [3].

In [2] we have proved that every n -dimensional real Bott manifold $M(A)$ admits an injective Seifert fibred structure which has the form $M(A) = T^k \times_{(\mathbb{Z}_2)^s} M(B)$, that is there is a k -torus action on $M(A)$ whose quotient space is an $(n-k)$ -dimensional real Bott orbifold $M(B)/(\mathbb{Z}_2)^s$ by some $(\mathbb{Z}_2)^s$ -action ($1 \leq s \leq k$). Moreover we have proved the smooth rigidity that two real Bott manifolds $M(A_i)$ $i = 1, 2$ are diffeomorphic if and only if the corresponding actions $((\mathbb{Z}_2)^{s_i}, M(B_i))$ are equivariantly diffeomorphic. When the low dimensional real Bott manifolds with $(\mathbb{Z}_2)^s$ -actions are classified, we can determine the diffeomorphism classes of higher dimensional ones by the above rigidity. We have classified real Bott manifolds until dimension 4 (see [4]).

The main purpose of this paper is to determine: (a) Diffeomorphism classes of 5-dimensional real Bott manifolds from the classifications of 2, 3, 4-dimensional ones with $(\mathbb{Z}_2)^s$ -actions ($s = 1, 2$), (b) Classification of certain type of n -dimensional real Bott manifolds $M(A)$.

We have obtained the following to (a), (Compare Theorem 3.10.)

Date: May 12, 2009.

2000 Mathematics Subject Classification. Primary 53C24; Secondary 57S25.

Key words and phrases. Bott tower, Bieberbach group, Flat Riemannian manifold, Real Bott manifold, Seifert fibration, Diffeomorphism.

Theorem A. There are 54 diffeomorphism classes of 5-dimensional real Bott manifolds.

Since each n -dimensional real Bott manifold has the form $M(A) = T^k \times_{(\mathbb{Z}_2)^s} M(B)$, it is shown that the diffeomorphism class of $M(A)$ is determined by the equivariant diffeomorphism class of the action $((\mathbb{Z}_2)^s, M(B))$ ($1 \leq s \leq k$) as above. We prove that if $k = 1$, there are 12-nonequivariant diffeomorphism classes of 4-dimensional real Bott manifolds with $\mathbb{Z}_2$ -actions. Then they create 29-diffeomorphism classes of such 5-dimensional real Bott manifolds. When $k = 2$, there are 4-nonequivariant diffeomorphism classes of 3-dimensional real Bott manifolds with $(\mathbb{Z}_2)^s$ -actions ($s = 1, 2$). Then from these, there are 19-diffeomorphism classes of the 5-dimensional real Bott manifolds. When $k = 3$, there are 2-nonequivariant diffeomorphism classes of 2-dimensional real Bott manifolds with $(\mathbb{Z}_2)^s$ -actions ($s = 1, 2$). Then there are 4-diffeomorphism classes of the 5-dimensional real Bott manifolds. When $k = 4$, the 1-dimensional real Bott manifold is S^1 with conjugate action of $\mathbb{Z}_2$, there exists only one such a 5-dimensional real Bott manifold. Finally if $k = 5$, the 5-dimensional real Bott manifold is T^5 . As a consequence, the total number of 5-dimensional diffeomorphism classes is 54. The details of the proof is in section 3.

It is far to determine the number of diffeomorphism classes of n -dimensional real Bott manifolds for $n \geq 6$. However, we shall solve the special types of higher dimensional real Bott manifolds.

Theorem B. Let T^k be the maximal torus action on an n -dimensional real Bott manifold ($n \geq 4$). If $k \geq n - 2$, then the diffeomorphism classes of such real Bott manifolds consists of 6.

See Corollary 4.2 for the proof.

Proposition C. The following hold.

- (i) Let $M(A)$ be a real Bott manifold which fibers S^1 over the real Bott manifold $M(B)$ for which $M(B)$ is either

$$T^k \times_{(\mathbb{Z}_2)^s} T^2 \text{ or } T^k \times_{(\mathbb{Z}_2)^s} K \text{ } (k \geq 2).$$

Here K is a Klein bottle. Then the diffeomorphism classes of such $M(A)$ is 3.

- (ii) Let $M(A)$ be a real Bott manifold which fibers S^1 over the real Bott manifold $M(B)$ where

$$M(B) = S^1 \times_{\mathbb{Z}_2} T^t \text{ } (t \geq 2),$$

then the diffeomorphism classes of such $M(A)$ is $[\frac{t}{2}] + 1$. Here $[x]$ is the Gauss integer.

- (iii) The diffeomorphism class is unique for the real Bott manifold of the form $M(A) = T^k \times_{\mathbb{Z}_2} T^{n-k}$ for any $n \geq 2$ and $k \geq 1$.

We prove these results in Section 4 (see Corollary 4.4, Corollary 4.5, Theorem 4.7, Proposition 4.6 respectively).

A special kind of Bott matrices is introduced in Section 1. We consider such a class of Bott matrices in (4.67).

Theorem D. Let $M(A) = S^1 \times_{\mathbb{Z}_2} M(B)$ be an n -dimensional real Bott manifold. Suppose that B is either one of the list in (4.67). Then $M(B)$ are diffeomorphic

each other and the number of diffeomorphism classes of such real Bott manifolds $M(A)$ above is $(k+1)2^{n-k-3}$ ($k \geq 2$, $n-k \geq 3$).

See Theorem 4.12 for the proof.

1. PRELIMINARIES

1.1. Seifert fiber space. Each i -th row of a Bott matrix A defines a $\mathbb{Z}_2$ -action on T^n by

$$g_i(z_1, \dots, z_n) = (z_1, \dots, z_{i-1}, -z_i, \tilde{z}_{i+1}, \dots, \tilde{z}_n), \quad (i = 1, \dots, n)$$

where (i, i) -(diagonal) entry 1 acts as $z_i \rightarrow -z_i$ while $\tilde{z}_j$ is either z_j or $\bar{z}_j$ depending on whether (i, j) -entry ($i < j$) is 0 or 1 respectively. Note that $\bar{z}$ is the conjugate of the complex number $z \in S^1$. It is always trivial; $z_j \rightarrow z_j$ whenever $j < i$. Here $(z_1, \dots, z_n)$ are the standard coordinates of the n -dimensional torus T^n whose universal covering is the n -dimensional euclidean space $\mathbb{R}^n$. The projection $p: \mathbb{R}^n \rightarrow T^n$ is denoted by

$$p(x_1, \dots, x_n) = (\exp(2\pi i x_1), \dots, \exp(2\pi i x_n)) = (z_1, \dots, z_n).$$

Those $\langle g_1, \dots, g_n \rangle$ constitute the generators of $(\mathbb{Z}_2)^n$. It is easy to see that $(\mathbb{Z}_2)^n$ acts freely on T^n such that the orbit space $M(A) = T^n / (\mathbb{Z}_2)^n$ is a smooth compact manifold. In this way, given a Bott matrix A of size n , we obtain a free action of $(\mathbb{Z}_2)^n$ on T^n .

Let $\pi(A) = \langle \tilde{g}_1, \dots, \tilde{g}_n \rangle$ be the lift of $(\mathbb{Z}_2)^n = \langle g_1, \dots, g_n \rangle$ to $\mathbb{R}^n$. Then we get

$$\tilde{g}_i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, \frac{1}{2} + x_i, \tilde{x}_{i+1}, \dots, \tilde{x}_n)$$

where $\tilde{x}_j$ is either x_j or $-x_j$. It is easy to see that $\pi(A)$ acts properly discontinuously and freely on $\mathbb{R}^n$ as euclidean motions. Note that $\pi(A)$ is a Bieberbach group which is a discrete uniform subgroup of the euclidean group $E(n) = \mathbb{R}^n \rtimes O(n)$ (cf.[5]). It follows that

$$M(A) = T^n / (\mathbb{Z}_2)^n = \mathbb{R}^n / \pi(A).$$

Now let us recall moves **I**, **II** and **III** [2] to a Bott matrix A of size n under which the diffeomorphism class of $M(A)$ does not change.

I. If the j -th column has all 0-entries except for the (j, j) -entry 1 for some $j > 1$, then interchange the j -th column and the $(j-1)$ -th column. Next, interchange the j -th row and the $(j-1)$ -th row.

This move **I** is interpreted in terms of the coordinates z_j 's of T^n and the generators g_j 's of $(\mathbb{Z}_2)^n$ as follows:

$$z_j \rightarrow z'_{j-1}, z_{j-1} \rightarrow z'_j, \quad g_j \rightarrow g'_{j-1}, g_{j-1} \rightarrow g'_j.$$

It is easy to see that the resulting matrix A' under move **I** is again a Bott matrix such that $M(A)$ is diffeomorphic to $M(A')$.

We perform move **I** iteratively to get a Bott matrix A'

$$(1.1) \quad A' = \left(\begin{array}{c|c} I_k & C \\ \hline 0 & B \end{array} \right) \quad B = \begin{pmatrix} 1 & & * \\ & \ddots & \\ & & 1 \end{pmatrix}$$

where I_k is a maximal block of identity matrix of size k , the entries of the $*$ are either 1 or 0, B is a Bott matrix of size $(n-k)$ which represents a real Bott manifold

$M(B) = T^{n-k}/(\mathbb{Z}_2)^{n-k}$. Since I_k is a maximal block of identity matrix, each $k+j$ ($j = 1, \dots, n-k$)-th column of A' has at least two non zero elements.

Associated with A' , the $(\mathbb{Z}_2)^n$ -action splits into $(\mathbb{Z}_2)^k \times (\mathbb{Z}_2)^{n-k}$ and T^n splits into $T^k \times T^{n-k}$. Hence

$$(1.2) \quad M(A) = T^n/(\mathbb{Z}_2)^n \cong \frac{T^k \times T^{n-k}}{(\mathbb{Z}_2)^k \times (\mathbb{Z}_2)^{n-k}} = T^k \times_{(\mathbb{Z}_2)^k} M(B) = M(A').$$

Note that the above $(\mathbb{Z}_2)^k$ -action of (1.2) is not necessarily effective on $M(B)$ but we can reduce it to the effective $(\mathbb{Z}_2)^s$ -action on $M(B)$ for some s ($1 \leq s \leq k$). In order to do so, we have two more moves.

II. If there is an m -th row ($1 \leq m \leq k$) whose entries in C are all zero, then divide $T^k \times M(B)$ by the corresponding $\mathbb{Z}_2$ -action. For example, suppose $M(A_1) = T^2 \times_{(\mathbb{Z}_2)^2} M(B)$ with

$$A_1 = \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

By move **II**, $M(A_1) \cong T^2 \times_{\mathbb{Z}_2} M(B)$.

III. If the p -th row and ℓ -th row ($1 \leq p < \ell \leq k$) have the common entries in C , then compose the $\mathbb{Z}_2$ -action of p -th row with ℓ -th row and divide $T^k \times M(B)$ by this $\mathbb{Z}_2$ -action. For example, suppose $M(A_2) = T^2 \times_{(\mathbb{Z}_2)^2} M(B)$ with

$$A_2 = \left(\begin{array}{cc|ccc} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

By move **III**, $M(A_2) \cong T^2 \times_{\mathbb{Z}_2} M(B) \cong M(A_1)$.

By an iteration of **II**, **III**, the quotient is again diffeomorphic to $T^k \times M(B)$ but eventually the $(\mathbb{Z}_2)^k$ -action is reduced to the effective $(\mathbb{Z}_2)^s$ -action on $T^k \times M(B)$. Therefore A' reduces to

$$(1.3) \quad A'' = \left(\begin{array}{c|c|c} I_{k-s} & 0 & 0 \\ \hline 0 & I_s & * \\ \hline 0 & 0 & B \end{array} \right)$$

in which

$$\begin{aligned} M(A') &= T^k \times_{(\mathbb{Z}_2)^k} M(B) \\ &= \frac{T^{k-s} \times T^s \times M(B)}{(\mathbb{Z}_2)^{k-s} \times (\mathbb{Z}_2)^s} = M(A''). \end{aligned}$$

Since $(\mathbb{Z}_2)^{k-s}$ acts trivially on $T^s \times M(B)$ then we have

$$M(A'') \cong T^k \times_{(\mathbb{Z}_2)^s} M(B).$$

From now on, we write $M(A)$ instead of $M(A'')$.

Remark 1.1. Since $(\mathbb{Z}_2)^s$ acts trivially on T^{k-s} ,

$$\begin{aligned} M(A) &\cong T^k \times_{(\mathbb{Z}_2)^s} M(B) = T^{k-s} \times T^s \times_{(\mathbb{Z}_2)^s} M(B) \\ &\cong (S^1)^{k-s} \times T^s \times_{(\mathbb{Z}_2)^s} M(B) \\ &= (S^1)^{k-s} \times M(B') \end{aligned}$$

where $M(B') = T^s \times_{(\mathbb{Z}_2)^s} M(B)$. That is, for $s < k$, a real Bott manifold $M(A)$ is the product of $(S^1)^{k-s}$ and an $(n-k+s)$ -dimensional real Bott manifold $M(B')$. In particular, if $M(A) = T^{n-1} \times_{\mathbb{Z}_2} S^1$ then it is diffeomorphic to $(S^1)^{n-2} \times$ Klein bottle.

Remark 1.2. From the submatrix $*$ of (1.3), the group $(\mathbb{Z}_2)^s = \langle g_{k-s+1}, \dots, g_k \rangle$ acts on $T^k \times M(B)$ by

$$\begin{aligned} (1.4) \quad g_i(z_1, \dots, z_{k-s+1}, \dots, z_k, [z_{k+1}, \dots, z_n]) \\ = (z_1, \dots, z_{k-s+1}, \dots, -z_i, \dots, z_k, [\tilde{z}_{k+1}, \dots, \tilde{z}_n]) \end{aligned}$$

where $\tilde{z} = \bar{z}$ or z . So there induces an action of $(\mathbb{Z}_2)^s$ on $M(B)$ by

$$(1.5) \quad g_i([z_{k+1}, \dots, z_n]) = [\tilde{z}_{k+1}, \dots, \tilde{z}_n].$$

Moreover in [2],

Theorem 1.3 (Structure). *Given a real Bott manifold $M(A)$, there exists a maximal T^k -action ($k \geq 1$) such that*

$$M(A) = T^k \times_{(\mathbb{Z}_2)^s} M(B)$$

is an injective Seifert fiber space over the $(n-k)$ -dimensional real Bott orbifold $M(B)/(\mathbb{Z}_2)^s$;

$$(1.6) \quad T^k \rightarrow M(A) \rightarrow M(B)/(\mathbb{Z}_2)^s.$$

There is a central extension of the fundamental group $\pi(A)$ of $M(A)$:

$$(1.7) \quad 1 \rightarrow \mathbb{Z}^k \rightarrow \pi(A) \rightarrow Q_B \rightarrow 1$$

such that

- (i) $\mathbb{Z}^k$ is the maximal central free abelian subgroup
- (ii) The induced group Q_B is the semidirect product $\pi(B) \rtimes (\mathbb{Z}_2)^s$ for which $\mathbb{R}^{n-k}/\pi(B) = M(B)$.

See [2] for the proof.

By this theorem, a real Bott manifold $M(A)$ which admits a maximal T^k -action ($k \geq 1$) can be created from an $(n-k)$ -dimensional real Bott manifold $M(B)$ by a $(\mathbb{Z}_2)^s$ -action, and the corresponding Bott matrix A has the form as in (1.3) above.

1.2. Affine maps between real Bott manifolds. Next, we can apply the following theorem to check whether two real Bott manifolds are diffeomorphic.

Theorem 1.4 (Rigidity). *Let $M(A_1)$, $M(A_2)$ be n -dimensional real Bott manifolds and $1 \rightarrow \mathbb{Z}^{k_i} \rightarrow \pi(A_i) \rightarrow Q_{B_i} \rightarrow 1$ be the associated group extensions ($i = 1, 2$). Then the following are equivalent:*

- (i) $\pi(A_1)$ is isomorphic to $\pi(A_2)$.
- (ii) There exists an isomorphism of $Q_{B_1} = \pi(B_1) \rtimes (\mathbb{Z}_2)^{s_1}$ onto $Q_{B_2} = \pi(B_2) \rtimes (\mathbb{Z}_2)^{s_2}$ preserving $\pi(B_1)$ and $\pi(B_2)$.
- (iii) The action $((\mathbb{Z}_2)^{s_1}, M(B_1))$ is equivariantly diffeomorphic to the action $((\mathbb{Z}_2)^{s_2}, M(B_2))$.

See [2] for the proof. Here Bott matrices A_1 and A_2 are created from B_1 and B_2 respectively.

Note that two real Bott manifolds $M(A_1)$ and $M(A_2)$ are diffeomorphic if and only if $\pi(A_1)$ is isomorphic to $\pi(A_2)$ by the Bieberbach theorem [5]. Moreover by Theorem 1.3 and 1.4 we have,

Remark 1.5. *Let real Bott manifolds $M(A_i) = T^{k_i} \times_{(\mathbb{Z}_2)^{s_i}} M(B_i)$ ($i = 1, 2$). If $M(A_1)$ and $M(A_2)$ are diffeomorphic then the following hold.*

- (i) $k_1 = k_2$.
- (ii) $M(B_1)$ and $M(B_2)$ are diffeomorphic.
- (iii) $s_1 = s_2$.

Therefore two real Bott manifolds which admit different maximal T^k -action are not diffeomorphic. If they have the same maximal T^k -action, then the quotients $((\mathbb{Z}_2)^{s_i}, M(B_i))$ are compared. If $M(B_1)$ is not diffeomorphic to $M(B_2)$ or $s_1 \neq s_2$, then $M(A_1)$ and $M(A_2)$ are not diffeomorphic. So our task is to distinguish the $(\mathbb{Z}_2)^{s_i}$ -action on $M(B_i)$ when it is the case that $s_1 = s_2 = s$ and $M(B_1)$ is diffeomorphic to $M(B_2)$.

1.3. Type of fixed point set. Note that from (1.5), the action of $(\mathbb{Z}_2)^s$ on $M(B)$ is defined by

$$\alpha[(z_1, \dots, z_{n-k})] = [\alpha(z_1, \dots, z_{n-k})] = [(\tilde{z}_1, \dots, \tilde{z}_{n-k})]$$

for $\alpha \in (\mathbb{Z}_2)^s$ and $\tilde{z} = z$ or $\bar{z}$. Since $M(B) = T^{n-k}/(\mathbb{Z}_2)^{n-k}$, the action $\langle \alpha \rangle$ lifts to a linear (affine) action on T^{n-k} naturally:

$$\alpha(z_1, \dots, z_{n-k}) = (\tilde{z}_1, \dots, \tilde{z}_{n-k}).$$

Then the fixed point set is characterized by the equation:

$$(\tilde{z}_1, \dots, \tilde{z}_{n-k}) = g(z_1, \dots, z_{n-k})$$

for some $g \in (\mathbb{Z}_2)^{n-k}$. It is also an affine subspace of T^{n-k} . So the fixed point sets of $(\mathbb{Z}_2)^s$ are affine subspaces in $M(B)$.

Let B be the Bott matrix as in (1.1). By a repetition of move **I**, B has the form

$$(1.8) \quad B = \begin{pmatrix} I_2 & C_{23} & \dots & \dots & C_{2\ell} \\ & I_3 & C_{34} & \dots & C_{3\ell} \\ & & \ddots & & \dots \\ & 0 & & I_{\ell-1} & C_{(\ell-1)\ell} \\ & & & & I_\ell \end{pmatrix}$$

where $\text{rank} B = n - k = \text{rank} I_2 + \dots + \text{rank} I_\ell$ and I_i ($i = 2, \dots, \ell$) is the identity matrix, C_{jt} ($j = 2, \dots, \ell - 1, t = 3, \dots, \ell$) is a $p_j \times q_t$ matrix ($p_j = \text{rank} I_j, q_t = \text{rank} I_t$).

Note that by the Bieberbach theorem (cf. [5]), if f is an isomorphism of $\pi(A_1)$ onto $\pi(A_2)$, then there exists an affine element $g = (h, H) \in A(n) = \mathbb{R}^n \rtimes \text{GL}(n, \mathbb{R})$ such that

$$(1.9) \quad f(r) = grg^{-1} \quad (\forall r \in \pi(A_1)).$$

Recall that if $M(A_1)$ is diffeomorphic to $M(A_2)$ then $M(B_1)$ is diffeomorphic to $M(B_2)$. This implies that B_1 and B_2 have the form as in (1.8).

Using (1.9) and according to the form of B in (1.8) we obtain that

$$(1.10) \quad g = \left(\begin{pmatrix} \mathbf{h}_1 \\ \mathbf{h}_2 \\ \vdots \\ \mathbf{h}_\ell \end{pmatrix}, \begin{pmatrix} H_1 & & & \\ & H_2 & & 0 \\ & & \ddots & \\ 0 & & & H_\ell \end{pmatrix} \right)$$

where $\mathbf{h}_i$ is an $s_i \times 1$ ($s_i = \text{rank} I_i$) column matrix ($\mathbf{h}_1$ is a $k \times 1$ column matrix), $H_i \in \text{GL}(s_i, \mathbb{R})$ ($i = 2, \dots, \ell$), $H_1 \in \text{GL}(k, \mathbb{R})$ (see Remark 3.2 [2]).

Let $\bar{f}: Q_{B_1} \rightarrow Q_{B_2}$ be the induced isomorphism from f (cf. Theorem 1.4). Now the affine equivalence $\bar{g}: \mathbb{R}^{n-k} \rightarrow \mathbb{R}^{n-k}$ has the form

$$(1.11) \quad \bar{g} = \left(\begin{pmatrix} \mathbf{h}_2 \\ \vdots \\ \mathbf{h}_\ell \end{pmatrix}, \begin{pmatrix} H_2 & & 0 \\ & \ddots & \\ 0 & & H_\ell \end{pmatrix} \right)$$

which is equivariant with respect to $\bar{f}$. The pair $(\bar{f}, \bar{g})$ induces an equivariant affine diffeomorphism $(\hat{f}, \hat{g}): ((\mathbb{Z}_2)^s, M(B_1)) \rightarrow ((\mathbb{Z}_2)^s, M(B_2))$.

Let $\text{rank} H_i = b_i$ ($i = 2, \dots, \ell$). (Note that $b_2 + \dots + b_\ell = n - k$.) Since $M(B_1) = T^{n-k}/(\mathbb{Z}_2)^{n-k}$, $\bar{g}$ induces an affine map $\tilde{g}$ of T^{n-k} . Put

$$X_{b_2} = \begin{pmatrix} x_1 \\ \vdots \\ x_{b_2} \end{pmatrix}, \dots, X_{b_\ell} = \begin{pmatrix} x_{b_{\ell'}+1} \\ \vdots \\ x_{b_{\ell'}+b_\ell} \end{pmatrix}, w_{b_i} = p(X_{b_i}) \in T^{b_i} \quad (i = 2, \dots, \ell),$$

$$b_{\ell'} = b_2 + \dots + b_{\ell-1}.$$

Since $\tilde{g}p = p\bar{g}$,

$$(1.12) \quad \tilde{g}({}^t w_{b_2}, \dots, {}^t w_{b_\ell}) = ({}^t w'_{b_2}, \dots, {}^t w'_{b_\ell})$$

where $w'_{b_i} = p(\mathbf{h}_i + H_i X_{b_i}) \in T^{b_i}$. That is, $\tilde{g}$ preserves each T^{b_i} of $T^{n-k} = T^{b_2} \times \dots \times T^{b_\ell}$, so does $\hat{g}$ on

$$M(B_1) = \{[z_1, \dots, z_{b_2}; z_{b_2+1}, \dots, z_{b_2+b_3}; \dots; z_{b_{\ell'}+1}, \dots, z_{b_{\ell'}+b_\ell}]\}.$$

We say that $\hat{g}$ preserves the type $(b_2, \dots, b_\ell)$ of $M(B_1)$. As $\hat{g}$ is $\hat{f}$ -equivariant, it also preserves the type corresponding to the fixed point sets between $((\mathbb{Z}_2)^s, M(B_1))$ and $((\mathbb{Z}_2)^s, M(B_2))$.

Proposition 1.6. *The $(\mathbb{Z}_2)^s$ -action on $M(B)$ is distinguished by the number of components and types of each positive dimensional fixed point subsets.*

See [2] for the proof.

Definition. We say that two Bott matrices A and A' are *equivalent* (denoted by $A \sim A'$) if $M(A)$ and $M(A')$ are diffeomorphic.

2. EXAMPLES

We shall give some real Bott manifolds in order to determine diffeomorphism classes of 5-dimensional ones. We introduce the following Bott matrices created

$$\text{from } B = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

$$A_1 = \left(\begin{array}{cc|cc} 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ \hline 0 & 0 & B \end{array} \right), \quad A_2 = \left(\begin{array}{cc|cc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & B \end{array} \right),$$

$$A_3 = \left(\begin{array}{cc|cc} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ \hline 0 & 0 & B \end{array} \right), \quad A_4 = \left(\begin{array}{cc|cc} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & B \end{array} \right).$$

Then we obtain the 5-dimensional real Bott manifolds $M(A_i)$ for which the $(\mathbb{Z}_2)^2$ -action on $M(B)$ is given by the first two rows of A_i ($i = 1, 2, 3, 4$). We prove that there are two distinct diffeomorphism classes among $M(A_i)$ ($i = 1, 2, 3, 4$).

- a) $M(A_1)$ is diffeomorphic to $M(A_2)$. For this, the $(\mathbb{Z}_2)^2$ -actions on $M(B)$ corresponding to A_1 and A_2 are given as follows:

$$(i) \quad g_1([z_3, z_4, z_5]) = [\bar{z}_3, \bar{z}_4, z_5] = [g_3(\bar{z}_3, \bar{z}_4, z_5)] = [-\bar{z}_3, z_4, z_5],$$

$$g_2([z_3, z_4, z_5]) = [z_3, \bar{z}_4, z_5],$$

$$(ii) \quad h_1([z_3, z_4, z_5]) = [z_3, \bar{z}_4, z_5], \quad h_2([z_3, z_4, z_5]) = [\bar{z}_3, z_4, z_5].$$

There is an equivariant diffeomorphism $\varphi: ((\mathbb{Z}_2)^2, M(B)) \rightarrow ((\mathbb{Z}_2)^2, M(B))$ defined by $\varphi([z_3, z_4, z_5]) = ([i z_3, z_4, z_5])$ such that $\varphi g_1 = h_2 \varphi$ and $\varphi g_2 = h_1 \varphi$. Hence the result follows from Theorem 1.4.

- b) $M(A_2)$ is not diffeomorphic to $M(A_3)$. If $M(A_2)$ and $M(A_3)$ are diffeomorphic, by Theorem 1.4 there is an equivariant diffeomorphism $\varphi: ((\mathbb{Z}_2)^2, M(B)) \rightarrow ((\mathbb{Z}_2)^2, M(B))$. Let $\bar{\varphi}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the lift of φ . According to the form of B , the affine element $\bar{\varphi}$ has the form

$$(2.1) \quad \bar{\varphi} = \left(\begin{pmatrix} a_2 \\ a_3 \\ a_4 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right)$$

for some $a_i \in \mathbb{R}$ ($i = 2, 3, 4$) (see (1.11)). Since $M(B) = T^3/(\mathbb{Z}_2)^3$, $\bar{\varphi}$ induces an affine map of T^3 . By the formula of (2.1), it preserves each S^1 of $T^3 = S^1 \times S^1 \times S^1$, so does φ on $M(B)$. Since φ is equivariant, it also preserves the type $(1, 1, 1)$ of the fixed point sets of $((\mathbb{Z}_2)^2, M(B))$. That is, if $[z_3, z_4, z_5]$ is a fixed point set of $((\mathbb{Z}_2)^2, M(B))$, then φ preserves each coordinate z_i ($i = 3, 4, 5$) (i.e., $\varphi[z_3, z_4, z_5] = [\exp(2\pi i a_2) z_3, \exp(2\pi i a_3) z_4, \exp(2\pi i a_4) z_5]$).

The fixed point sets of $((\mathbb{Z}_2)^2, M(B))$ corresponding to A_2 and A_3 are as follows:

- (i) 3 components $T^2 = \{[z_3, 1, z_5], [1, z_4, z_5], [\mathbf{i}, z_4, z_5]\}$,
 4 components $S^1 = \{[z_3, \mathbf{i}, 1], [\mathbf{i}, 1, z_5], [z_3, \mathbf{i}, \mathbf{i}], [1, 1, z_5]\}$,
 4 points $\{[\mathbf{i}, \mathbf{i}, 1], [\mathbf{i}, \mathbf{i}, \mathbf{i}], [1, \mathbf{i}, 1], [1, \mathbf{i}, \mathbf{i}]\}$,
- (ii) 3 components $T^2 = \{[z_3, z_4, \mathbf{i}], [1, z_4, z_5], [z_3, z_4, 1]\}$,
 4 components $S^1 = \{[1, z_4, 1], [\mathbf{i}, 1, z_5], [1, z_4, \mathbf{i}], [\mathbf{i}, \mathbf{i}, z_5]\}$,
 and 4 points $\{[\mathbf{i}, \mathbf{i}, 1], [\mathbf{i}, \mathbf{i}, \mathbf{i}], [\mathbf{i}, 1, 1], [\mathbf{i}, 1, \mathbf{i}]\}$.

We see that the number of components of fixed point sets of $((\mathbb{Z}_2)^2, M(B))$ corresponding to A_2 and A_3 is the same. Since the type of fixed point set is preserved, φ maps $T^2 = \{[z_3, 1, z_5]\}$, $z_3, z_5 \in S^1$ ((i) in A_2) onto the fixed point set $T^2 = \{[w_3, \exp(2\pi i a_3), w_5]\}$ ($w_3, w_5 \in S^1$) of A_3 . However there is no type of such fixed point set in (ii) of A_3 . Therefore by Proposition 1.6, $M(A_2)$ and $M(A_3)$ are not diffeomorphic.

- c) $M(A_3)$ is diffeomorphic to $M(A_4)$. In this case, the $(\mathbb{Z}_2)^2$ -actions on $M(B)$ corresponding to A_3 and A_4 are given as follows:

- (i) $g_1([z_3, z_4, z_5]) = [\bar{z}_3, z_4, z_5]$, $g_2([z_3, z_4, z_5]) = [z_3, z_4, \bar{z}_5]$,
- (ii) $h_1([z_3, z_4, z_5]) = [\bar{z}_3, z_4, \bar{z}_5]$, $h_2([z_3, z_4, z_5]) = [\bar{z}_3, z_4, z_5]$.

We change the generator h_1 by h'_1 :

$$h'_1([z_3, z_4, z_5]) = h_1 h_2 [z_3, z_4, z_5] = [z_3, z_4, \bar{z}_5].$$

Define an equivariant diffeomorphism $\varphi: ((\mathbb{Z}_2)^2, M(B)) \rightarrow ((\mathbb{Z}_2)^2, M(B))$ to be $\varphi([z_3, z_4, z_5]) = ([z_3, z_4, z_5])$ such that $\varphi g_1 = h_2 \varphi$ and $\varphi g_2 = h'_1 \varphi$. Hence $M(A_3)$ is diffeomorphic to $M(A_4)$ by Theorem 1.4.

3. FIVE-DIMENSIONAL REAL BOTT MANIFOLDS

Before giving the classification of 5-dimensional real Bott manifolds, we recall the classification of 2, 3, 4-dimensional ones as stated in [3], [4].

Theorem 3.1. *The diffeomorphism classes of 2-dimensional real Bott manifolds consist of two. The corresponding Bott matrices are as follows.*

$$A_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Theorem 3.2. *The diffeomorphism classes of 3-dimensional real Bott manifolds consist of four. The corresponding Bott matrices are classified into four equivalence classes as follows:*

- a) $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$
- b) $\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$
- c) $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$
- d) $I_3.$

Theorem 3.3. *The diffeomorphism classes of 4-dimensional real Bott manifolds consist of twelve. The corresponding Bott matrices are classified into twelve equivalence classes as follows:*

[illegible]

$$\begin{aligned}
\text{ix)} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
\text{x)} & \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
\text{xi)} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
\text{xii)} & I_4.
\end{aligned}$$

Using the classification results of Theorem 3.1, 3.2, 3.3, we shall classify 5-dimensional real Bott manifolds.

3.1. S^1 -actions with 4-dimensional quotients.

The Bott matrices of $M(A)$ admitting S^1 -actions have the following form

$$\left(\begin{array}{c|cccc} 1 & 1 & a_{13} & a_{14} & a_{15} \\ \hline 0 & B & & & \end{array} \right)$$

where $a_{13}, a_{14}, a_{15} \in \{0, 1\}$. In this case $M(B)$ corresponds to the Bott matrices B in Theorem 3.3. Taking the first Bott matrix from i) as B , we consider the following Bott matrices.

$$A_1 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_2 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 1 & 0 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_3 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 1 & 1 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

Then the fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_1 , A_2 and A_3 respectively are as follows: (1) T^3 , T^2 , 4 points, (2) 2 components T^2 , 4 components S^1 , (3) T^2 , 4 components S^1 , 4 points. (1), (2) and (3) have the different fixed point sets each other so each A_i ($i = 1, 2, 3$) is not equivalent by Proposition 1.6.

The following Bott matrices are created from the first Bott matrix in ii).

$$A_4 = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_5 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_6 = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 1 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_4 , A_5 and A_6 are obtained as: (1) 3 components T^2 , 4 points, (2) T^3 , 4 components S^1 , (3) 8 components S^1 . In view of the fixed points, similarly A_i ($i = 4, 5, 6$) are not equivalent to each other.

The following Bott matrices are created from the first Bott matrix in iii).

$$A_7 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_8 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 1 & 0 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$A_9 = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 1 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{10} = \left(\begin{array}{c|cccc} 1 & 1 & 0 & 1 & 1 \\ \hline 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_7 , A_8 , A_9 and A_{10} are as follows:

- (1) T^3 , T^2 , S^1 , 2 points,
- (2) 2 components $T^2 = \{[1, z_3, 1, z_5], [\mathbf{i}, \mathbf{i}, z_4, z_5]\}$, 3 components $S^1 = \{[\mathbf{i}, 1, 1, z_5], [1, z_3, \mathbf{i}, 1], [1, z_3, \mathbf{i}, \mathbf{i}]\}$, 2 points= $\{[\mathbf{i}, 1, \mathbf{i}, 1], [\mathbf{i}, 1, \mathbf{i}, \mathbf{i}]\}$,
- (3) 2 components $T^2 = \{[1, z_3, z_4, 1], [1, z_3, z_4, \mathbf{i}]\}$, 3 components $S^1 = \{[\mathbf{i}, \mathbf{i}, \mathbf{i}, z_5], [\mathbf{i}, 1, z_4, 1], [\mathbf{i}, 1, z_4, \mathbf{i}]\}$, 2 points= $\{[\mathbf{i}, \mathbf{i}, 1, 1], [\mathbf{i}, \mathbf{i}, 1, \mathbf{i}]\}$,
- (4) T^2 , 5 components S^1 , 2 points.

Note that the fixed point sets of (2) and (3) coincide, but the type of them are different. (Compare **b**) in Section 2 for the type (1,1,1,1).) Hence A_8 and A_9 are not equivalent. As the fixed point sets (1), (4) and (2) (or (3)) are all different, each A_i ($i = 7, 8, 9, 10$) is not equivalent.

The following Bott matrices are created from the first Bott matrix in iv).

$$\begin{aligned}
A_{11} &= \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{12} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 1 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{13} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 0 & 1 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{14} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 1 & 1 \\ \hline 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).
\end{aligned}$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_{11} , A_{12} , A_{13} and A_{14} are as follows: (1) T^3 , 3 components S^1 , 2 points, (2) T^2 , 5 components S^1 , 2 points, (3) 3 components T^2 , S^1 , 2 points, (4) 2 components T^2 , 3 components S^1 , 2 points. By Proposition 1.6, A_i ($i = 11, 12, 13, 14$) are not equivalent to each other.

The Bott matrices A_i ($i = 15, 16, 17, 18$) below are created from the first Bott matrix in v) while A_{19} is created from the second Bott matrix in v).

$$\begin{aligned}
A_{15} &= \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{16} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 1 & 0 \\ \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{17} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 0 & 1 \\ \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{18} &= \left(\begin{array}{c|ccccc} 1 & 1 & 0 & 1 & 1 \\ \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{19} &= \left(\begin{array}{c|ccccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).
\end{aligned}$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_{15} , A_{16} , A_{17} , A_{18} and A_{19} are as follows: (1) T^3 , 2 components S^1 , 4 points, (2) 3 components T^2 , 2 components S^1 , (3) 2 components T^2 , 2 components S^1 , 4 point, (4) T^2 , 6 components S^1 , (5) 2 components T^2 , 4 components S^1 . By Proposition 1.6, A_i ($i = 15, 16, 17, 18, 19$) are not equivalent to each other.

The following Bott matrices are created from the Bott matrix vi).

$$\begin{aligned}
A_{20} &= \left(\begin{array}{c|cccc} 1 & 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{21} &= \left(\begin{array}{c|ccccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).
\end{aligned}$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_{20} and A_{21} are as follows: (1) T^3 , 8 points, (2) 2 components T^2 , 4 components S^1 . By Proposition 1.6, A_{20} and A_{21} are not equivalent.

The Bott matrix

$$A_{22} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

is created from the first Bott matrix in vii).

The following Bott matrices are created from the first Bott matrix in viii).

$$A_{23} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{24} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 1 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_{23} and A_{24} are as follows: (1) 2 components T^2 , 8 points, (2) 8 components S^1 . By Proposition 1.6, A_{23} is not equivalent to A_{24} .

The following Bott matrices are created from the first Bott matrix in ix).

$$A_{25} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{26} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 1 \\ \hline 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

The fixed point sets of the $\mathbb{Z}_2$ -actions on $M(B)$ corresponding to A_{25} and A_{26} are as follows: (1) 2 components T^2 , 2 components S^1 , 4 points, (2) 6 components S^1 , 4 points. By Proposition 1.6, A_{25} and A_{26} are not equivalent.

The Bott matrix A_{27} (resp. A_{28}) below is created from the first Bott matrix in x) (resp. xi)).

$$A_{27} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{28} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 1 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

Finally from I_4 , we get $A_{29} = \left(\begin{array}{c|cccc} 1 & 1 & 1 & 1 & 1 \\ \hline 0 & I_4 \end{array} \right).$

Since each Bott matrix B of i) to xii) is not equivalent to each other, the resulting Bott matrix A is not equivalent. Totally, 29 Bott matrices A_i ($i = 1, \dots, 29$) are not equivalent to each other. When we take the second Bott matrix B' from i), the resulting Bott matrix A' gives an action $(\mathbb{Z}_2, M(B'))$. We can check that $(\mathbb{Z}_2, M(B'))$ is equivariantly diffeomorphic to one of the actions $(\mathbb{Z}_2, M(B))$ corresponding to A_1, A_2, A_3 by the ad hoc argument. (Compare Section 2 for the argument to find an equivariant diffeomorphism.) Once there exists such an equivariant diffeomorphism, A' is equivalent to one of A_1, A_2, A_3 by Theorem 1.4.

Similarly, if A' is another Bott matrix created from the first Bott matrix in i), we can check that the corresponding $(\mathbb{Z}_2, M(B))$ is equivariantly diffeomorphic to one of the actions $(\mathbb{Z}_2, M(B))$ corresponding to A_1, A_2, A_3 . (Note that the total number of Bott matrices created from the first Bott matrix in i) is 8.) This argument works not only the case i) but also the cases from ii) to xii). As a consequence the Bott matrix A' created from Bott matrices from ii) to xii) is equivalent to one of A_i 's ($i = 4, \dots, 29$). In summary, we obtain the following but the proof is omitted because of a tedious argument.

Lemma 3.4. *A Bott matrix created from any one of Bott matrices of Theorem 3.3 is equivalent to one of the Bott matrices A_i ($i = 1, \dots, 29$) above.*

Proposition 3.5. *There are 29 diffeomorphism classes of the case S^1 -actions with 4-dimensional quotients.*

3.2. T^2 -actions with 3-dimensional quotients.

The Bott matrices of $M(A)$ admitting T^2 -actions have the following form

$$\left(\begin{array}{c|c} I_2 & * \\ \hline 0 & B \end{array} \right).$$

The following Bott matrices are created from the first Bott matrix B of a) in Theorem 3.2.

$$\begin{aligned} A_{30} &= \left(\begin{array}{c|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{31} &= \left(\begin{array}{c|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\ A_{32} &= \left(\begin{array}{c|ccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{33} &= \left(\begin{array}{c|ccc} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\ A_{34} &= \left(\begin{array}{c|ccc} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{35} &= \left(\begin{array}{c|ccc} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right). \end{aligned}$$

The fixed point sets of the $(\mathbb{Z}_2)^s$ -actions ($s = 1, 2$) on $M(B)$ corresponding to A_{30} , A_{31} , A_{32} , A_{33} , A_{34} and A_{35} are as follows: (1) T^2 , S^1 , 2 points, (2) 3 components S^1 , 2 points, (3) 3 components T^2 , 4 components S^1 , 4 points, (4) T^2 , 8 components S^1 , 4 points, (5) 3 components T^2 , 4 components S^1 , 4 points, (6) 2 components T^2 , 6 components S^1 , 4 points. Compared (3) with (5), we see from **b)** in Section 2 that A_{32} is not equivalent to A_{34} . By Proposition 1.6, Bott matrices A_i ($i = 30, 31$) (resp. A_j ($j = 32, 33, 34, 35$)) are not equivalent to each other. Moreover, by Remark 1.5, Bott matrices A_i ($i = 30, 31$) are not equivalent to A_j ($j = 32, 33, 34, 35$) because the $(\mathbb{Z}_2)^2$ -action corresponding to A_j ($j = 32, 33, 34, 35$) cannot be reduced to a $\mathbb{Z}_2$ -action.

The following Bott matrices are created from the Bott matrix b) in Theorem 3.2.

$$\begin{aligned}
A_{36} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{37} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ \hline 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{38} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{39} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{40} &= \left(\begin{array}{cc|ccc} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \\ \hline 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).
\end{aligned}$$

The fixed point sets of the $(\mathbb{Z}_2)^s$ -actions ($s = 1, 2$) on $M(B)$ corresponding to A_{36} , A_{37} , A_{38} , A_{39} and A_{40} are as follows: (1) T^2 , 4 points, (2) 4 components S^1 , (3) 2 components T^2 , 4 components S^1 , 8 points, (4) 3 components T^2 , 4 components S^1 , 4 points, (5) 12 components S^1 . By Remark 1.5, Bott matrices A_i ($i = 36, 37$) are not equivalent to A_j ($j = 38, 39, 40$) because the $(\mathbb{Z}_2)^2$ -action corresponding to A_j ($j = 38, 39, 40$) cannot be reduced to a $\mathbb{Z}_2$ -action. On the other hand, by Proposition 1.6, Bott matrices A_i ($i = 36, 37$) (resp. A_j ($j = 38, 39, 40$)) are not equivalent to each other.

The Bott matrices A_i ($i = 41, 42, 43, 44$) below are created from the first Bott matrix in c) of Theorem 3.2 while A_{45} is created from the second Bott matrix in c).

$$\begin{aligned}
A_{41} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ \hline 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{42} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{43} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ \hline 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) & A_{44} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \\ \hline 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \\
A_{45} &= \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).
\end{aligned}$$

The fixed point sets of the $(\mathbb{Z}_2)^2$ -actions on $M(B)$ corresponding to A_{42} , A_{43} , A_{44} and A_{45} are as follows: (1) 3 components T^2 , 4 components S^1 , 4 points, (2) T^2 , 8 components S^1 , 4 points, (3) 2 components T^2 , 4 components S^1 , 8 points,

(4) 2 components T^2 , 6 components S^1 , 4 points. By Remark 1.5, A_{41} is not equivalent to A_i ($i = 42, 43, 44, 45$) because the $(\mathbb{Z}_2)^2$ -action corresponding to A_i ($i = 42, 43, 44, 45$) cannot be reduced to a $\mathbb{Z}_2$ -action. Then by Proposition 1.6, Bott matrices A_i ($i = 42, 43, 44, 45$) are not equivalent to each other.

The following Bott matrices are created from I_3 .

$$A_{46} = \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & I_3 & & \end{array} \right) \quad A_{47} = \left(\begin{array}{cc|ccc} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & I_3 & & \end{array} \right)$$

$$A_{48} = \left(\begin{array}{cc|ccc} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & I_3 & & \end{array} \right).$$

The fixed point sets of the $(\mathbb{Z}_2)^2$ -actions on $M(B)$ corresponding to A_{47} and A_{48} are as follows: (1) 2 components T^2 , 4 components S^1 , 8 points, (2) 12 components S^1 . By Remark 1.5, A_{46} is not equivalent to A_i ($i = 47, 48$), and by Proposition 1.6, A_{47} is not equivalent to A_{48} .

Since each Bott matrix B of a) to d) is not equivalent to each other, the resulting Bott matrix A is not equivalent. Totally, 19 Bott matrices A_i ($i = 30, \dots, 48$) are not equivalent to each other.

When we take the second Bott matrix B' from a) of Theorem 3.2, the resulting Bott matrix A' gives an action $((\mathbb{Z}_2)^s, M(B'))$ ($s = 1, 2$). We can check that $((\mathbb{Z}_2)^s, M(B'))$ is equivariantly diffeomorphic to one of the actions $((\mathbb{Z}_2)^s, M(B))$ corresponding to A_i ($i = 30, \dots, 35$) by the ad hoc argument. (Compare Section 2 for the argument to find an equivariant diffeomorphism.) Once there exists such an equivariant diffeomorphism, A' is equivalent to one of A_i 's ($i = 30, \dots, 35$) by Theorem 1.4. Similarly, if A' is another Bott matrix created from the first Bott matrix in a) of Theorem 3.2, we can check that the corresponding $((\mathbb{Z}_2)^s, M(B))$ ($s = 1, 2$) is equivariantly diffeomorphic to one of the actions $((\mathbb{Z}_2)^s, M(B))$ corresponding to A_i ($i = 30, \dots, 35$). (Note that the total number of Bott matrices created from the first Bott matrix in a) is 76.) This argument also works for the case b), c) and d). As a consequence the Bott matrix A' created from Bott matrices in b), c) and d) is equivalent to one of A_i 's ($i = 36, \dots, 48$). In summary, we obtain the following.

Lemma 3.6. *A Bott matrix created from any one of Bott matrices of Theorem 3.2 is equivalent to one of the Bott matrices A_i ($i = 30, \dots, 48$) above.*

Proposition 3.7. *There are 19 diffeomorphism classes of the case T^2 -actions with 3-dimensional quotients.*

3.3. T^3 -actions with 2-dimensional quotients.

The Bott matrices of $M(A)$ admitting T^3 -actions have the following form

$$\left(\begin{array}{c|c} I_3 & * \\ \hline 0 & B \end{array} \right).$$

In this case a Bott matrix B is either A_1 or A_2 in Theorem 3.1. The Bott matrices A_{49} and A_{50} (resp. A_{51} and A_{52}) below are created from A_1 (resp. A_2).

$$A_{49} = \left(\begin{array}{ccc|cc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ \hline 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{50} = \left(\begin{array}{ccc|cc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ \hline 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$A_{51} = \left(\begin{array}{ccc|cc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ \hline 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \quad A_{52} = \left(\begin{array}{ccc|cc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ \hline 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

Since A_1 and A_2 in Theorem 3.1 are not equivalent, none of A_{49} and A_{50} is equivalent to A_{51} or A_{52} . Then A_{49} (resp. A_{51}) is not equivalent to A_{50} (resp. A_{52}), because $(\mathbb{Z}_2)^2$ -action on $M(B)$ corresponding to A_{50} (or A_{52}) cannot be reduced to a $\mathbb{Z}_2$ -action. If A' is another Bott matrix created from A_1 in Theorem 3.1, we can check that the corresponding $((\mathbb{Z}_2)^s, M(A_1))$ ($s = 1, 2$) is equivariantly diffeomorphic to one of the actions $((\mathbb{Z}_2)^s, M(A_1))$ corresponding to A_{49} and A_{50} by the ad hoc argument. Once there exists such an equivariant diffeomorphism, A' is equivalent to A_{49} or A_{50} by Theorem 1.4. This argument works also for the case A_2 in Theorem 3.1. As a consequence another Bott matrix A' created from A_2 is equivalent to A_{51} or A_{52} . Thus we obtain the following.

Lemma 3.8. *A Bott matrix created from any one of Bott matrices in Theorem 3.1 is equivalent to one of the Bott matrices A_i ($i = 49, 50, 51, 52$) above.*

Proposition 3.9. *There are 4 diffeomorphism classes of the case T^3 -actions with 2-dimensional quotients.*

3.4. T^4 -actions with one-dimensional quotients.

The Bott matrices of $M(A)$ admitting T^4 -actions have the following form

$$\left(\begin{array}{cccc|c} I_4 & & & & * \\ \hline 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

In this case $M(B) = M(1) = S^1$. It is easy to check by using moves **II** and **III**, it consists of just one diffeomorphism class, where the corresponding Bott matrix is

$$A_{53} = \left(\begin{array}{cccc|c} I_4 & & & & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

Obviously the corresponding Bott matrix of size 5 of a real Bott manifold admitting T^5 -action is the identity matrix of rank 5. Combined with Proposition 3.5, 3.7, 3.9 and the case of T^4 -actions above we get the following theorem.

Theorem 3.10. *The diffeomorphism classes of 5-dimensional real Bott manifolds consist of 54.*

4. CLASSIFICATION OF n -DIMENSIONAL REAL BOTT MANIFOLDS

In this section we shall prove some results regarding the classification of certain types of n -dimensional real Bott manifolds.

Theorem 4.1. *The number of diffeomorphism classes of n -dimensional real Bott manifolds ($n \geq 4$) which admit the maximal T^{n-2} -actions (i.e. $s = 1, 2$) is 4:*

$$M(A) = T^{(n-2)} \times_{(\mathbb{Z}_2)^s} M(B).$$

Proof. Since $M(B)$ is a 2-dimensional real Bott manifold, the real Bott manifolds $M(A)$ created from $M(B)$ correspond to the following Bott matrices

$$(4.1) \quad \left(\begin{array}{c|c} I_{n-2} & * \\ \hline 0 & B_1 \end{array} \right),$$

$$(4.2) \quad \left(\begin{array}{c|c} I_{n-2} & * \\ \hline 0 & B_2 \end{array} \right)$$

where $B_1 = I_2$, $B_2 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Let us consider (4.1). If the entries in each row of $*$ are the same then by moves **II** or **III**, (4.1) is equivalent to

$$(4.3) \quad \left(\begin{array}{c|cc} I_{n-2} & 0 & 0 \\ \hline & 1 & 1 \\ & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).$$

Otherwise by moves **II**, **III** or the equivariant diffeomorphism $\varphi: ((\mathbb{Z}_2)^2, M(B_1)) \rightarrow ((\mathbb{Z}_2)^2, M(B_1))$ defined by $\varphi[z_{n-1}, z_n] = [z_{n-1}, z_n]$, (4.1) is equivalent to

$$(4.4) \quad \left(\begin{array}{c|cc} I_{n-2} & 0 & 0 \\ \hline & 1 & 0 \\ & 0 & 1 \\ 0 & 1 & 0 \\ & 0 & 1 \end{array} \right).$$

However (4.3) is not equivalent to (4.4) because the $(\mathbb{Z}_2)^2$ -action on $M(B_1)$ corresponding to (4.4) cannot be reduced to a $\mathbb{Z}_2$ -action on it.

Let us consider (4.2). If the entries in each row of $*$ are the same then (4.2) is equivalent to

$$(4.5) \quad \left(\begin{array}{c|cc} I_{n-2} & 0 & 0 \\ \hline & 1 & 1 \\ & 1 & 1 \\ 0 & 0 & 1 \end{array} \right)$$

or if the entries in the second column of $*$ are all zero then (4.2) is equivalent to

$$(4.6) \quad \left(\begin{array}{c|cc} I_{n-2} & 0 & 0 \\ \hline & 1 & 0 \\ & 1 & 1 \\ 0 & 0 & 1 \end{array} \right)$$

by moves **II** or **III**. However (4.5) and (4.6) are equivalent by the equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B_2)) \rightarrow (\mathbb{Z}_2, M(B_2))$ defined by $\varphi([z_{n-1}, z_n]) = [\mathbf{i}z_{n-1}, z_n]$.

Otherwise (4.2) is equivalent to

$$(4.7) \quad \left(\begin{array}{c|cc} & 0 & 0 \\ I_{n-2} & 1 & 0 \\ \hline & 0 & 1 \\ 0 & 1 & 1 \\ \hline & 0 & 1 \end{array} \right)$$

by moves **II**, **III** or the equivariant diffeomorphism $\varphi: ((\mathbb{Z}_2)^2, M(B_2)) \rightarrow ((\mathbb{Z}_2)^2, M(B_2))$ defined by $\varphi[z_{n-1}, z_n] = [z_{n-1}, z_n]$. Moreover (4.6) and (4.7) are not equivalent because the $(\mathbb{Z}_2)^2$ -action on $M(B_2)$ corresponding to (4.7) cannot be reduced to a $\mathbb{Z}_2$ -action on it.

Therefore Bott matrices (4.1) and (4.2) give 4 distinct diffeomorphism classes of real Bott manifolds $M(A)$. $\square$

Corollary 4.2. *For any $n \geq 4$, the number of diffeomorphism classes of n -dimensional real Bott manifolds which admit the maximal T^k -actions ($k = n-2, n-1, n$) is 6.*

Proof. It is easy to check that real Bott manifolds $M(A_1) = T^{n-1} \times_{\mathbb{Z}_2} S^1$ and $M(A_2) = T^n$ have only one diffeomorphism class respectively, because the corresponding Bott matrices are $A_1 = \left(\begin{array}{c|cc} I_{n-1} & * & \\ \hline 0 & 1 & 1 \end{array} \right)$ and $A_2 = I_n$ respectively. More-

over, A_1 reduces to $A'_1 = \left(\begin{array}{c|cc} I_{n-1} & 0 & \\ \hline 0 & 1 & 1 \end{array} \right)$ by move **III**. $\square$

Corollary 4.3. *If $M(A) = S^1 \times_{\mathbb{Z}_2} M(B)$ where $M(B) = T^k \times_{\mathbb{Z}_2} S^1$, then for any $k \geq 1$ there is only one diffeomorphism class.*

Proof. Since $M(B) = T^k \times_{\mathbb{Z}_2} S^1$, as shown in the proof of Corollary 4.2,

$$B = \left(\begin{array}{ccc|cc} & I_k & & * & \\ & & & 1 & \\ 0 & \dots & 0 & 1 & 1 \end{array} \right).$$

The Bott matrices A created from B are

$$\left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 1 \\ \hline 0 & B & & & \end{array} \right) \text{ and } \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 \\ \hline 0 & B & & & \end{array} \right)$$

which are equivalent by the equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B)) \rightarrow (\mathbb{Z}_2, M(B))$ defined by $\varphi([z_2, \dots, z_{k+1}, z_{k+2}]) = [z_2, \dots, \mathbf{i}z_{k+1}, z_{k+2}]$. $\square$

Corollary 4.4. *For any $k \geq 2$, there are 3 diffeomorphism classes in $(k+3)$ -dimensional real Bott manifolds $M(A) = S^1 \times_{\mathbb{Z}_2} M(B)$ where $M(B) = T^k \times_{(\mathbb{Z}_2)^s} T^2$ ($s = 1, 2$).*

Proof. Since $M(B) = T^k \times_{(\mathbb{Z}_2)^s} T^2$ ($s = 1, 2$), there are 2 distinct diffeomorphism classes of $M(B)$ which correspond to the following Bott matrices

$$(4.8) \quad B_1 = \left(\begin{array}{c|cc} I_k & 0 & 0 \\ \hline & 1 & 1 \\ & 1 & 0 \\ 0 & 0 & 1 \end{array} \right), \quad B_2 = \left(\begin{array}{c|cc} I_k & 0 & 0 \\ \hline & 1 & 0 \\ & 0 & 1 \\ 0 & 1 & 0 \\ & 0 & 1 \end{array} \right)$$

(see (4.3) and (4.4)).

The Bott matrices of size $(k+3)$ created from (4.8) with the $\mathbb{Z}_2$ -actions are as follows

$$A_1 = \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & * & * \\ \hline 0 & & & B_1 & & \end{array} \right), \quad A_2 = \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & * & * \\ \hline 0 & & & B_2 & & \end{array} \right).$$

The following Bott matrices in (4.9) (resp. (4.10)) derived from A_1

$$(4.9) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 0 \\ \hline 0 & & & B_1 & & \end{array} \right), \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 1 & 1 \\ \hline 0 & & & B_1 & & \end{array} \right)$$

$$(4.10) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 1 & 0 \\ \hline 0 & & & B_1 & & \end{array} \right), \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 1 \\ \hline 0 & & & B_1 & & \end{array} \right)$$

are equivalent by the equivariant diffeomorphism

$$\begin{aligned} \varphi: (\mathbb{Z}_2, M(B_1)) &\rightarrow (\mathbb{Z}_2, M(B_1)) \\ \varphi([z_2, \dots, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, \dots, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}]. \end{aligned}$$

Moreover, Bott matrices in (4.9) are not equivalent to (4.10) because the maximal fixed point sets of $(\mathbb{Z}_2, M(B_1))$ corresponding to the Bott matrices in (4.9) and (4.10) are T^2 and S^1 respectively.

It is easy to see that each Bott matrix derived from A_2 is equivalent to

$$(4.11) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 0 \\ \hline 0 & & & B_2 & & \end{array} \right)$$

by $\varphi: (\mathbb{Z}_2, M(B_2)) \rightarrow (\mathbb{Z}_2, M(B_2))$ which is defined by one of the following

$$\begin{aligned} \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, \dots, \mathbf{i}z_k, z_{k+1}, z_{k+2}, z_{k+3}], \\ \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, z_3, \dots, z_k, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}], \\ \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, z_3, \dots, \mathbf{i}z_k, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}]. \end{aligned}$$

Obviously, (4.11) is not equivalent to the Bott matrices in (4.9) and (4.10) because they are created from two nonequivalent Bott matrices in (4.8). Therefore there are 3 equivalence classes of the Bott matrices corresponding to $M(A)$. $\square$

Corollary 4.5. *For any $k \geq 2$, there are 3 diffeomorphism classes in $(k+3)$ -dimensional real Bott manifolds $M(A) = S^1 \times_{\mathbb{Z}_2} M(B)$ where $M(B) = T^k \times_{(\mathbb{Z}_2)^s} K$, (K =Klein bottle, $s = 1, 2$).*

Proof. Since $M(B) = T^k \times_{(\mathbb{Z}_2)^s} K$ ($s = 1, 2$), there are 2 distinct diffeomorphism classes of $M(B)$ corresponding to the following Bott matrices (see (4.6) and (4.7))

$$(4.12) \quad B_1 = \left(\begin{array}{c|cc} I_k & 0 & 0 \\ \hline & 1 & 0 \\ & 1 & 1 \\ 0 & 0 & 1 \end{array} \right), B_2 = \left(\begin{array}{c|cc} & 0 & 0 \\ I_k & 1 & 0 \\ \hline & 0 & 1 \\ & 1 & 1 \\ 0 & 0 & 1 \end{array} \right).$$

The Bott matrices of size $(k+3)$ created from (4.12) with the $\mathbb{Z}_2$ -actions are as follows

$$A_1 = \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & * & * \\ \hline 0 & & & B_1 & & \end{array} \right), A_2 = \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & * & * \\ \hline 0 & & & B_2 & & \end{array} \right).$$

The following Bott matrices in (4.13) (resp. (4.14)) derived from A_1

$$(4.13) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 0 \\ \hline 0 & & & B_1 & & \end{array} \right), \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 1 & 0 \\ \hline 0 & & & B_1 & & \end{array} \right)$$

$$(4.14) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 1 \\ \hline 0 & & & B_1 & & \end{array} \right), \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 1 & 1 \\ \hline 0 & & & B_1 & & \end{array} \right)$$

are equivalent by the equivariant diffeomorphism

$$\begin{aligned} \varphi: (\mathbb{Z}_2, M(B_1)) &\rightarrow (\mathbb{Z}_2, M(B_1)) \\ \varphi([z_2, \dots, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, \dots, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}]. \end{aligned}$$

On the other hand, Bott matrices in (4.13) are not equivalent to (4.14) because the maximal fixed point sets of $(\mathbb{Z}_2, M(B))$ corresponding to the Bott matrices in (4.13) and (4.14) are T^2 and S^1 respectively.

It is easy to see that each Bott matrix derived from A_2 is equivalent to

$$(4.15) \quad \left(\begin{array}{c|cccc} 1 & 1 & \dots & 1 & 0 & 0 \\ \hline 0 & & & B_2 & & \end{array} \right)$$

by $\varphi: (\mathbb{Z}_2, M(B_2)) \rightarrow (\mathbb{Z}_2, M(B_2))$ which is defined by one of the following

$$\begin{aligned} \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, \dots, \mathbf{i}z_k, z_{k+1}, z_{k+2}, z_{k+3}], \\ \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, z_3, \dots, z_k, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}], \\ \varphi([z_2, \dots, z_k, z_{k+1}, z_{k+2}, z_{k+3}]) &= [z_2, z_3, \dots, \mathbf{i}z_k, \mathbf{i}z_{k+1}, z_{k+2}, z_{k+3}]. \end{aligned}$$

Obviously, the Bott matrix (4.15) is not equivalent to the Bott matrices in (4.13) and (4.14) because they are created from two nonequivalent Bott matrices in (4.12). Therefore there are 3 equivalence classes of the Bott matrices corresponding to $M(A)$. $\square$

Proposition 4.6. *If $M(A) = T^k \times_{\mathbb{Z}_2} T^{n-k}$, then for any $n \geq 2$ and $k \geq 1$ there is only one diffeomorphism class.*

Proof. Since $M(A)$ admits the maximal T^k -action and A is created from I_{n-k} , there is only one the Bott matrix A , namely

$$(4.16) \quad A = \left(\begin{array}{c|ccc} I_k & & 0 & \\ \hline & 1 & \dots & 1 \\ 0 & & I_{n-k} & \end{array} \right).$$

$\square$

Now if we create Bott matrices from (4.16) (for $k = 1$) with $\mathbb{Z}_2$ -actions then we will get a classification of the corresponding real Bott manifolds as follows.

Theorem 4.7. *For any $k \geq 2$, there are $\lfloor \frac{k}{2} \rfloor + 1$ diffeomorphism classes in $(k+2)$ -dimensional real Bott manifolds $M(A_i) = S^1 \times_{\mathbb{Z}_2} M(B)$ ($i = 1, \dots, 2^k$), where $M(B) = S^1 \times_{\mathbb{Z}_2} T^k$. Here $[x]$ is the Gauss integer.*

Proof. Since $M(B) = S^1 \times_{\mathbb{Z}_2} T^k$,

$$(4.17) \quad B = \left(\begin{array}{c|ccc} 1 & 1 & \dots & 1 \\ \hline 0 & & I_k & \end{array} \right).$$

The Bott matrices A_i of size $(k+2)$ created from (4.17) with $\mathbb{Z}_2$ -actions are as follows

$$(4.18) \quad A_i = \left(\begin{array}{c|ccc} 1 & 1 & * & \\ \hline 0 & 1 & \dots & 1 \\ & & I_k & \end{array} \right) = \left(\begin{array}{c|ccc} 1 & 1 & * & \\ \hline 0 & & B & \end{array} \right), \quad (i = 1, \dots, 2^k).$$

We apply the different $\mathbb{Z}_2$ -actions on $M(B)$ such that the Bott matrices A_i are as follows

$$(4.19) \quad \begin{aligned} A_1 &= \left(\begin{array}{c|ccc} 1 & \{1\}_2 & \{0\}_3 & \dots & \{0\}_{2+k} \\ \hline 0 & & B & & \end{array} \right), \\ A_2 &= \left(\begin{array}{c|ccc} 1 & \{1\}_2 & \{1\}_3 & \{0\}_4 & \dots & \{0\}_{2+k} \\ \hline 0 & & B & & & \end{array} \right), \\ &\vdots \\ A_{\lfloor \frac{k}{2} \rfloor + 1} &= \left(\begin{array}{c|ccccccc} 1 & \{1\}_2 & \dots & \{1\}_{1+(\lfloor \frac{k}{2} \rfloor + 1)} & \{0\}_{2+(\lfloor \frac{k}{2} \rfloor + 1)} & \dots & \{0\}_{2+k} \\ \hline 0 & & & B & & & \end{array} \right). \end{aligned}$$

It is easy to check that the maximal fixed point sets of $(\mathbb{Z}_2, M(B))$ corresponding to A_i ($i = 1, 2, \dots, \lfloor \frac{k}{2} \rfloor + 1$) are $T^k, T^{k-1}, \dots, T^{k-\lfloor \frac{k}{2} \rfloor}$ respectively. Hence they are not equivalent to each other. Here $\{y\}_i$ means y in the i -th spot.

On the other hand, for $\lfloor \frac{k}{2} \rfloor + 1 < l \leq (k+1)$, Bott matrix

$$(4.20) \quad \left(\begin{array}{c|ccccccc} 1 & \{1\}_2 & \dots & \{1\}_{1+l} & \{0\}_{2+l} & \dots & \{0\}_{2+k} \\ \hline 0 & & & B & & & \end{array} \right)$$

is equivalent to one of the Bott matrices in (4.19). To show this, consider the g_1 -action corresponding to (4.20):

$$\begin{aligned} g_1([z_2, \dots, z_{1+l}, z_{l+2}, \dots, z_{k+2}]) &= [\bar{z}_2, \dots, \bar{z}_{l+1}, \overbrace{z_{l+2}, \dots, z_{k+2}}^{(k+1)-l}] \\ &= [g_2(\bar{z}_2, \dots, \bar{z}_{l+1}, z_{l+2}, \dots, z_{k+2})] \\ &= [-\bar{z}_2, z_3, \dots, z_{l+1}, \bar{z}_{l+2}, \dots, \bar{z}_{k+2}]. \end{aligned}$$

Since $(k+1) - l < k - \lfloor \frac{k}{2} \rfloor \leq \lfloor \frac{k}{2} \rfloor + 1$, there is an equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B)) \rightarrow (\mathbb{Z}_2, M(B))$ defined by $\varphi([z_2, \dots, z_{l+1}, z_{l+2}, \dots, z_{k+2}]) = [\mathbf{i}z_2, z_{l+2}, \dots, z_{k+2}, z_3, \dots, z_{l+1}]$ such that $\varphi g_1 = h_1 \varphi$ for some h_1 -action corresponding to one of the Bott matrices in (4.19).

The other Bott matrices A_i ($i \neq 1, \dots, \lfloor \frac{k}{2} \rfloor + 1$) may have the form

$$(4.21) \quad A' = \left(\begin{array}{c|ccc} 1 & 1 & \hat{1} & \dots & \hat{1} \\ \hline 0 & & B & & \end{array} \right),$$

where $\hat{1} \in \{0, 1\}$. If the number of entries $\hat{1}$ for $\hat{1} = 1$ is less than or equal to $\lfloor \frac{k}{2} \rfloor$ then A' is equivalent to one of the Bott matrices in (4.19), otherwise A' is equivalent to (4.20). We shall prove it in the following way. Suppose that the number of entries $\hat{1}$ for $\hat{1} = 1$ is t . Applying move **I** on A' such that the entries $\hat{1}$ for $\hat{1} = 1$ are placed in series, we get a new Bott matrix

$$A'' = \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{1\}_3 \quad \dots \quad \{1\}_{2+t} \quad \{0\}_{3+t} \quad \dots \quad \{0\}_{2+k}}{B} \right),$$

which is still equivalent to A' . Obviously, $A'' = A_i$ for some $i = 1, \dots, \lfloor \frac{k}{2} \rfloor + 1$ if $0 \leq t \leq \lfloor \frac{k}{2} \rfloor$, or A'' is the same as (4.20) if $t > \lfloor \frac{k}{2} \rfloor$. Hence A' is equivalent to one of the Bott matrices in (4.19). This completes the proof of theorem. $\square$

From now on, we use the notation $(\mathbb{Z}_2, M(B_j))_i$ which means that the $\mathbb{Z}_2$ -action on $M(B_j)$ corresponds to a Bott matrix A_{ij} .

Lemma 4.8. *Let $M(A_{i1}) = S^1 \times_{\mathbb{Z}_2} M(B_1)$ ($i = 1, \dots, n-2$) be n -dimensional real Bott manifolds creating from an $(n-1)$ -dimensional real Bott manifold $M(B_1)$. Such real Bott manifolds $M(A_{i1})$ corresponding to A_{i1} in (4.22) are not diffeomorphic to each other.*

(4.22)

$$\begin{aligned} A_{11} &= \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{0\}_3 \quad \dots \quad \{0\}_n}{B_1} \right), \\ A_{21} &= \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{1\}_3 \quad \{0\}_4 \quad \dots \quad \{0\}_n}{B_1} \right), \\ &\vdots \\ A_{(n-k-1)1} &= \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{1\}_3 \quad \dots \quad \{1\}_{1+(n-k-1)} \quad \{0\}_{1+(n-k)} \quad \dots \quad \{0\}_n}{B_1} \right), \\ &\vdots \\ A_{(n-2)1} &= \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{1\}_3 \quad \dots \quad \{1\}_{1+(n-2)} \quad \{0\}_n}{B_1} \right), \end{aligned}$$

where

$$(4.23) \quad B_1 = \begin{pmatrix} 1 & 1 & 1 & \dots & \dots & \dots & 1 \\ & 1 & 1 & \dots & \dots & \dots & 1 \\ & & \ddots & & & & \vdots \\ & & & 1 & 1 & \dots & 1 \\ 0 & & & & & I_k & \end{pmatrix},$$

$k \geq 2$, and $n - k \geq 3$.

Proof. Recall that, if $M(A_{m1})$ is diffeomorphic to $M(A_{q1})$ (i.e., A_{m1} is equivalent to A_{q1} , $m \neq q$), by Theorem 1.4, there is an equivariant diffeomorphism

$$\begin{aligned} (\Phi, \varphi): (\mathbb{Z}_2 = \langle \alpha \rangle, M(B_1))_m &\rightarrow (\mathbb{Z}_2 = \langle \beta \rangle, M(B_1))_q, \text{ such that} \\ \varphi(\alpha[z_2, \dots, z_n]) &= \Phi(\alpha)\varphi[z_2, \dots, z_n] = \beta\varphi[z_2, \dots, z_n]. \end{aligned}$$

Let $\bar{\varphi}: \mathbb{R}^{n-1} \rightarrow \mathbb{R}^{n-1}$ be the lift of φ . According to the form of B_1 , the affine element $\bar{\varphi}$ has the form

$$(4.24) \quad \bar{\varphi} = \left(\left(\begin{array}{c} \mathbf{a} \\ \hline \mathbf{b} \end{array} \right), \left(\begin{array}{ccc|c} 1 & & 0 & \\ & \ddots & & 0 \\ 0 & & 1 & \\ \hline & 0 & & D \end{array} \right) \right)$$

where D is a nonsingular submatrix of rank k , ${}^t\mathbf{a} = (a_2, \dots, a_{n-k})$ and ${}^t\mathbf{b} = (b_{n-k+1}, \dots, b_n)$ (see (1.11)). Since $M(B_1) = T^{n-1}/(\mathbb{Z}_2)^{n-1}$, $\bar{\varphi}$ induces an affine map $\tilde{\varphi}$ of T^{n-1} .

$$(4.25) \quad \text{Put } X = \begin{pmatrix} x_{n-k+1} \\ \vdots \\ x_n \end{pmatrix}. \text{ Since } \tilde{\varphi}p = p\bar{\varphi},$$

$$\begin{aligned} \tilde{\varphi}(z_2, \dots, z_{n-k}, z_{n-k+1}, \dots, z_n) &= (\ell_2 z_2, \dots, \ell_{n-k} z_{n-k}, p^t(\mathbf{b} + DX)) \\ &= (\ell_2 z_2, \dots, \ell_{n-k} z_{n-k}, c_{n-k+1} w_{n-k+1}, \dots, c_n w_n) \end{aligned}$$

where $\ell_p = \exp(2\pi i a_p)$ ($p = 2, \dots, n-k$), $c_s = \exp(2\pi i b_s)$ ($s = n-k+1, \dots, n$), $(w_{n-k+1}, \dots, w_n) = p^t(DX)$.

On the other hand, since $M(B_1) = T^{n-1}/(\mathbb{Z}_2)^{n-1}$, the action $\langle \alpha \rangle$ lifts to an action on T^{n-1} such that we have the commutative diagram

$$\begin{array}{ccc} (\mathbb{Z}_2)^{n-1} & & (\mathbb{Z}_2)^{n-1} \\ \downarrow & & \downarrow \\ (\alpha, T^{n-1}) & \xrightarrow{\tilde{\varphi}} & (g\beta, T^{n-1}) \\ \text{\scriptsize } Pr \downarrow & & \text{\scriptsize } Pr \downarrow \\ (\alpha, M(B_1))_m & \xrightarrow{\varphi} & (\beta, M(B_1))_q \end{array}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$. This means that

$$\begin{aligned} Pr(\tilde{\varphi}(\alpha(z_2, \dots, z_n))) &= \varphi(Pr(\alpha(z_2, \dots, z_n))) = \varphi(\alpha(Pr(z_2, \dots, z_n))) \\ &= \Phi(\alpha)\varphi(Pr(z_2, \dots, z_n)) = \beta\varphi(Pr(z_2, \dots, z_n)) \\ &= \beta Pr(\tilde{\varphi}(z_2, \dots, z_n)) = Pr(\beta\tilde{\varphi}(z_2, \dots, z_n)), \end{aligned}$$

(i.e.,

$$(4.26) \quad \tilde{\varphi}(\alpha(z_2, \dots, z_n)) = g\beta\tilde{\varphi}(z_2, \dots, z_n).)$$

Note that g_i ($i = 2, \dots, n$) corresponds to the i -th row of A_{m1} and A_{q1} . This implies that $\tilde{\varphi}$ maps the fixed point set of (α, T^{n-1}) to that of $(g\beta, T^{n-1})$ diffeomorphically. From the commutative diagram, we also have, for $g \in (\mathbb{Z}_2)^{n-1}$,

$$\begin{aligned} Pr(\tilde{\varphi}(g(z_2, \dots, z_n))) &= \varphi(Pr(g(z_2, \dots, z_n))) = \varphi(Pr(z_2, \dots, z_n)) \\ &= Pr(\tilde{\varphi}(z_2, \dots, z_n)). \end{aligned}$$

Hence there is an element $h \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.27) \quad \begin{aligned} \tilde{\varphi}(g(z_2, \dots, z_n)) &= h\tilde{\varphi}(z_2, \dots, z_n) \\ \tilde{\varphi}g\tilde{\varphi}^{-1} &= h. \end{aligned}$$

Recall from (4.22), that

$$A_{i1} = \left(\frac{1}{0} \middle| \begin{array}{cccccc} \{1\}_2 & \{1\}_3 & \dots & \{1\}_{1+i} & \{0\}_{2+i} & \dots & \{0\}_n \end{array} \right),$$

for $i = 1, \dots, n-2$.

To show that Bott matrices A_{i1} ($i = 1, \dots, n-2$) are not equivalent to each other, we shall prove the following cases.

a). Bott matrices A_{i1} ($i = 1, \dots, n-k-1$) are not equivalent to each other.

Suppose that A_{l1} is equivalent to A_{p1} where $1 \leq l < p \leq n-k-1$ ($l = 1, \dots, n-k-2$; $p = 2, \dots, n-k-1$). By the definition, the α -action and β -action on T^{n-1} corresponding to A_{l1} and A_{p1} are as follows

$$(4.28) \quad \begin{aligned} \alpha(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{l+1}, z_{l+2}, \dots, z_n), \\ \beta(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{l+1}, \bar{z}_{l+2}, \dots, \bar{z}_{p+1}, z_{p+2}, \dots, z_n). \end{aligned}$$

Then from (4.26), we have

$$(4.29) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (\ell_2\bar{z}_2, \dots, \ell_{l+1}\bar{z}_{l+1}, \ell_{l+2}z_{l+2}, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \dots, \overline{\ell_{l+1}z_{l+1}}, \overline{\ell_{l+2}z_{l+2}}, \dots, \overline{\ell_{p+1}z_{p+1}}, \ell_{p+2}z_{p+2}, \dots, \ell_{n-k}z_{n-k}, \\ &\quad c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \end{aligned}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$. Since

$$(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) = g(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n),$$

g is a composition of an even number of generators $\{g_2, \dots, g_{n-k}\}$.

On the other hand, since $g_t(z_t) = -z_t$ ($t = 2, \dots, n-k$) and

$$(4.30) \quad (\ell_{l+2}z_{l+2}, \dots, \ell_{n-k}z_{n-k}) = g(\overline{\ell_{l+2}z_{l+2}}, \dots, \overline{\ell_{p+1}z_{p+1}}, \ell_{p+2}z_{p+2}, \dots, \ell_{n-k}z_{n-k}),$$

$g \notin \langle g_{l+2}, \dots, g_{n-k} \rangle$. So, the last possibility is that $g \in \langle g_2, \dots, g_{l+1} \rangle$. If this is the case, and since g is a composition of an even number of generators $\{g_2, \dots, g_{l+1}\}$, it contradicts (4.30).

b). Bott matrices A_{i1} ($i = n-k, \dots, n-2$) are not equivalent to each other.

Suppose that $A_{(n-k+l)1}$ is equivalent to $A_{(n-k+p)1}$ where $n-k \leq n-k+l < n-k+p \leq n-2$ ($l = 0, \dots, k-3$; $p = 1, \dots, k-2$). By the definition, the α -action and β -action on T^{n-1} corresponding to $A_{(n-k+l)1}$ and $A_{(n-k+p)1}$ are as follows

$$(4.31) \quad \begin{aligned} \alpha(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+l+1}, z_{n-k+l+2}, \dots, z_n), \\ \beta(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+p+1}, z_{n-k+p+2}, \dots, z_n). \end{aligned}$$

Then from (4.26), we have

$$\begin{aligned}
 (4.32) \quad & \tilde{\varphi}(\alpha(z_2, \dots, z_n)) = g\beta\tilde{\varphi}(z_2, \dots, z_n), \\
 & (\ell_2 \bar{z}_2, \dots, \ell_{n-k} \bar{z}_{n-k}, c_{n-k+1} u'_{n-k+1}, \dots, c_n u'_n) \\
 & = g(\overline{\ell_2 z_2}, \dots, \overline{\ell_{n-k} z_{n-k}}, \overline{c_{n-k+1} w_{n-k+1}}, \dots, \overline{c_{n-k+p+1} w_{n-k+p+1}}, \\
 & \quad c_{n-k+p+2} w_{n-k+p+2}, \dots, c_n w_n)
 \end{aligned}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$, where

$$(u'_{n-k+1}, \dots, u'_n) = p((-x_{n-k+1}, \dots, -x_{n-k+l+1}, x_{n-k+l+2}, \dots, x_n)^t D).$$

Now we consider the following cases for g .

b1). If $g = g_t g'$ with $g_t \in \{g_2, \dots, g_{n-k-1}\}$, $g' \in \langle g_{t+1}, \dots, g_n \rangle$ then

$$\ell_{t+1} \bar{z}_{t+1} = g(\overline{\ell_{t+1} z_{t+1}}) = \begin{cases} -\ell_{t+1} z_{t+1} & \text{if } g = g_t g_{t+1} g'', g'' \in \langle g_{t+2}, \dots, g_n \rangle \\ \ell_{t+1} z_{t+1} & \text{if } g = g_t g''. \end{cases}$$

Hence we get a contradiction. That is, such $g = g_t g'$ cannot occur.

b2). If $g = g_{n-k} \hat{g}$ where $\hat{g} \in \langle g_{n-k+1}, \dots, g_n \rangle$, then

$$\ell_{n-k} \bar{z}_{n-k} = g(\overline{\ell_{n-k} z_{n-k}}) = -\overline{\ell_{n-k} z_{n-k}}.$$

This implies that $\ell_{n-k} = \pm \mathbf{i}$. Therefore

$$\begin{aligned}
 \tilde{\varphi}(z_2, \dots, z_n) &= (\ell_2 z_2, \dots, \ell_{n-k-1} z_{n-k-1}, \pm \mathbf{i} z_{n-k}, p^t(\mathbf{b} + DX)), \\
 \tilde{\varphi}^{-1}(z_2, \dots, z_n) &= (\bar{\ell}_2 \bar{z}_2, \dots, \bar{\ell}_{n-k-1} \bar{z}_{n-k-1}, \mp \mathbf{i} z_{n-k}, p^t(-D^{-1} \mathbf{b} + D^{-1} X)).
 \end{aligned}$$

Now, from (4.27), we consider

$$\begin{aligned}
 & \tilde{\varphi} g_2 \tilde{\varphi}^{-1}(z_2, \dots, z_n) \\
 &= (-z_2, \ell_3^2 \bar{z}_3, \dots, \ell_{n-k-1}^2 \bar{z}_{n-k-1}, -\bar{z}_{n-k}, \exp(4\pi \mathbf{i} b_{n-k+1}) \bar{z}_{n-k+1}, \dots, \exp(4\pi \mathbf{i} b_n) \bar{z}_n) \\
 &= g_2(z_2, \overline{\ell_3^2 z_3}, \dots, \overline{\ell_{n-k-1}^2 z_{n-k-1}}, -z_{n-k}, \\
 & \quad \exp(4\pi(-\mathbf{i}) b_{n-k+1}) z_{n-k+1}, \dots, \exp(4\pi(-\mathbf{i}) b_n) z_n) \\
 &= g_2 h(z_2, \dots, z_n)
 \end{aligned}$$

where

$$\begin{aligned}
 (4.33) \quad h(z_2, \dots, z_n) &= (z_2, \overline{\ell_3^2 z_3}, \dots, \overline{\ell_{n-k-1}^2 z_{n-k-1}}, -z_{n-k}, \\
 & \quad \exp(4\pi(-\mathbf{i}) b_{n-k+1}) z_{n-k+1}, \dots, \exp(4\pi(-\mathbf{i}) b_n) z_n).
 \end{aligned}$$

We shall check that $h \notin \langle g_3, \dots, g_n \rangle$ (i.e., $g_2 h \notin (\mathbb{Z}_2)^{n-1}$).

Suppose that $h \in \langle g_3, \dots, g_n \rangle$. Since $h(z_{n-k}) = -z_{n-k}$ in (4.33), we may write $h = \hat{h} g_{n-k} h''$, where $\hat{h}$ is a composition of an even number of generators $\{g_3, \dots, g_{n-k-1}\}$, $h'' \in \langle g_{n-k+1}, \dots, g_n \rangle$. However such h implies that $h(z_{n-k+1}, \dots, z_n) = (\pm \bar{z}_{n-k+1}, \dots, \pm \bar{z}_n)$. This contradicts (4.33). Similarly for $h = g_{n-k} h''$. Thus $g_2 h \notin (\mathbb{Z}_2)^{n-1}$. That is, such $g = g_{n-k} \hat{g}$ cannot occur.

b3). If $g = \hat{g}$ satisfies (4.32), then from (4.31),

$$\begin{aligned}
 & \hat{g}\beta(z_2, \dots, z_n) = \\
 & (\bar{z}_2, \dots, \bar{z}_{n-k}, \pm \bar{z}_{n-k+1}, \dots, \pm \bar{z}_{n-k+l+1}, \dots, \pm \bar{z}_{n-k+p+1}, \pm z_{n-k+p+2}, \dots, \pm z_n).
 \end{aligned}$$

Then we obtain that the fixed point set of (α, T^{n-1}) is

$$Fix \alpha = (\{\pm 1\}_2, \dots, \{\pm 1\}_{n-k+l+1}, z_{n-k+l+2}, \dots, z_n)$$

and the fixed point set of $(\hat{g}\beta, T^{n-1})$ is

$$Fix \hat{g}\beta = (\{\pm 1\}_2, \dots, \{\pm 1\}_{n-k}, \{\star\}_{n-k+1}, \dots, \{\star\}_{n-k+p+1}, z_{n-k+p+2}, \dots, z_n)$$

with $\star \in \{\pm 1, \pm i\}$. Then by (4.26), we have

$$\begin{aligned} dim(Fix \alpha) &= dim(Fix g\beta) \\ k - l - 1 &= k - p - 1. \end{aligned}$$

Hence we get a contradiction. That is, such $g = \hat{g}$ cannot occur.

c). Each A_{i1} ($i = 1, \dots, n - k - 1$) is not equivalent to each $A_{(n-k+j)1}$ ($j = 0, \dots, k - 2$).

Suppose that A_{i1} is equivalent to $A_{(n-k+j)1}$. By the definition, the α -action and β -action on T^{n-1} corresponding to A_{i1} and $A_{(n-k+j)1}$ are as follows

$$(4.34) \quad \begin{aligned} \alpha(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{i+1}, z_{i+2}, \dots, z_{n-k}, \dots, z_n), \\ \beta(z_2, \dots, z_n) &= (\bar{z}_2, \dots, \bar{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+j+1}, z_{n-k+j+2}, \dots, z_n). \end{aligned}$$

Then by (4.26), we have

$$(4.35) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (\ell_2\bar{z}_2, \dots, \ell_{i+1}\bar{z}_{i+1}, \ell_{i+2}z_{i+2}, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \dots, \overline{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j+1}w_{n-k+j+1}}, \\ &\quad c_{n-k+j+2}w_{n-k+j+2}, \dots, c_nw_n). \end{aligned}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$.

On the other hand, since

$$\begin{aligned} (c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) &= \\ g(\overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j+1}w_{n-k+j+1}}, c_{n-k+j+2}w_{n-k+j+2}, \dots, c_nw_n), \end{aligned}$$

there is no $g \in \langle g_2, \dots, g_n \rangle$ satisfying (4.35). This completes the proof of Lemma. $\square$

Lemma 4.9. Let $M(A_{j'1}) = S^1 \times_{\mathbb{Z}_2} M(B_1)$ ($j = 1, \dots, k, (k+1)$) be n -dimensional real Bott manifolds creating from an $(n-1)$ -dimensional real Bott manifold $M(B_1)$. Such real Bott manifolds $M(A_{j'1})$ corresponding to Bott matrices derived from $A_{j'1}$ ($j = 1, \dots, k, (k+1)$) in (4.36) are not diffeomorphic to each other.

(4.36)

$$A_{(1')1} = \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{0\}_3 \quad \{\hat{1}\}_4 \quad \cdots \quad \{\hat{1}\}_{(n-k)} \quad \{0\} \quad \cdots \quad \{0\}_{(n-k)+k}}{B_1} \right),$$

$$A_{(2')1} = \left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{0\}_3 \quad \{\hat{1}\}_4 \quad \cdots \quad \{\hat{1}\}_{(n-k)} \quad \{1\}_{n-k+1} \quad \{0\} \quad \cdots \quad \{0\}_n}{B_1} \right),$$

⋮

 $A_{(k')1} =$

$$\left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{0\}_3 \quad \{\hat{1}\}_4 \quad \cdots \quad \{\hat{1}\}_{(n-k)} \quad \{1\}_{n-k+1} \quad \cdots \quad \{1\}_{n-1} \quad \{0\}_n}{B_1} \right),$$

 $A_{(k+1)')1} =$

$$\left(\frac{1}{0} \middle| \frac{\{1\}_2 \quad \{0\}_3 \quad \{\hat{1}\}_4 \quad \cdots \quad \{\hat{1}\}_{(n-k)} \quad \{1\}_{n-k+1} \quad \cdots \quad \{1\}_{n-1} \quad \{1\}_n}{B_1} \right)$$

where $\hat{1}$ is either 0 or 1,

$$(4.37) \quad (\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) \neq (0, \dots, 0)$$

(resp. $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) \neq \overbrace{(0, \dots, 0)}^l, 1, \dots, 1)$, $l = 0, 1, \dots, n-k-3$)

for Bott matrix $A_{j'1}$ ($j = 1, \dots, k$) (resp. $A_{(k+1)')1}$), the Bott matrix B_1 is as in (4.23), $k \geq 2$, and $n-k \geq 3$. (That is, there are $k(2^{n-k-3}-1) + (2^{n-k-3} - (n-k-2))$ nonequivalent Bott matrices derived from (4.36).)

Proof. For brevity, we can write Bott matrices (4.36) in this way

 $A_{j'1} =$

$$\left(\frac{1}{0} \middle| \frac{1 \quad 0 \quad \hat{1} \quad \cdots \quad \{\hat{1}\}_{n-k} \quad \{1\}_{n-k+1} \quad \cdots \quad \{1\}_{n-k+(j-1)} \quad \{0\} \quad \cdots \quad \{0\}_n}{B_1} \right)$$

for $j = 1, \dots, k, (k+1)$.

To show that Bott matrices derived from $A_{j'1}$ ($j = 1, \dots, k, (k+1)$) in (4.36) are not equivalent to each other, we shall prove the following claims by using the argument at the beginning of the proof of Lemma (4.8).

Claim 1. Bott matrices $A_{j'1}$ ($j = 1, \dots, k, (k+1)$) are not equivalent to each other.

Suppose that $A_{l'1}$ is equivalent to $A_{p'1}$ where $1 \leq l < p \leq k+1$ ($l = 1, \dots, k$; $p = 2, \dots, k+1$). By the definition, the α -action and β -action on T^{n-1} corresponding to $A_{l'1}$ and $A_{p'1}$ are as follows

$$(4.38) \quad \alpha(z_2, \dots, z_n) = (\bar{z}_2, z_3, \widehat{\bar{z}_4}^\alpha, \dots, \widehat{z_{n-k}}^\alpha, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+l-1}, z_{n-k+l}, \dots, z_n),$$

$$\beta(z_2, \dots, z_n) = (\bar{z}_2, z_3, \widehat{\bar{z}_4}^\beta, \dots, \widehat{z_{n-k}}^\beta, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+l-1}, \dots, \bar{z}_{n-k+p-1}, z_{n-k+p}, \dots, z_n).$$

Here $\hat{z}_j^\alpha (\in \{z_j, \bar{z}_j\})$ means an α -action on z_j . Similarly for $\hat{z}_j^\beta$. Note that $\hat{z}_j$ is either z_j or $\bar{z}_j$ depending on whether $\hat{1}$ is 0 or 1 respectively. Then by (4.26), we have

$$\begin{aligned}
 (4.39) \quad & \tilde{\varphi}(\alpha(z_2, \dots, z_n)) = g\beta\tilde{\varphi}(z_2, \dots, z_n), \\
 & (\ell_2\bar{z}_2, \ell_3z_3, \ell_4\hat{z}_4^\alpha, \dots, \ell_{n-k}\hat{z}_{n-k}^\alpha, c_{n-k+1}w'_{n-k+1}, \dots, c_nw'_n) \\
 & = g(\overline{\ell_2z_2}, \ell_3z_3, \overline{\ell_4z_4}^\beta, \dots, \overline{\ell_{n-k}z_{n-k}}^\beta, \\
 & \quad \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+p-1}w_{n-k+p-1}}, c_{n-k+p}w_{n-k+p}, \dots, c_nw_n),
 \end{aligned}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$, where

$$(w'_{n-k+1}, \dots, w'_n) = p((-x_{n-k+1}, \dots, -x_{n-k+l-1}, x_{n-k+l}, \dots, x_n)^t D).$$

Obviously, $g \in \langle g_2, g_3 \rangle$ does not satisfy (4.39), because it implies that

$$(4.40) \quad \ell_3z_3 = g(\ell_3z_3) = \begin{cases} \overline{\ell_3z_3} & \text{if } g = g_2 \\ -\ell_3z_3 & \text{if } g = g_3 \\ -\overline{\ell_3z_3} & \text{if } g = g_2g_3. \end{cases}$$

Next we consider the following cases for $g \in \langle g_4, \dots, g_n \rangle$.

Case 1. Let $g = g_t g'$ where $g_t \in \{g_4, \dots, g_{n-k}\}$ ($t = 4, \dots, n-k$), $g' \in \langle g_{t+1}, \dots, g_n \rangle$.

Note that since $g_t(z_t) = -z_t$, $g(z_t) = -z_t$. If $\hat{z}_t^\alpha \neq \hat{z}_t^\beta$ (resp. $\hat{z}_t^\alpha = \hat{z}_t^\beta = z_t$) then $g = g_t g'$ does not satisfy (4.39), because it implies that $\ell_t z_t = g(\ell_t z_t) = -\overline{\ell_t z_t}$ (resp. $\ell_t z_t = g(\ell_t z_t) = -\ell_t z_t$). If $\hat{z}_t^\alpha = \hat{z}_t^\beta = \bar{z}_t$ then $\ell_t \bar{z}_t = g(\ell_t \bar{z}_t) = -\overline{\ell_t \bar{z}_t}$. This implies that $\ell_t = \pm \mathbf{i}$. Therefore

$$\begin{aligned}
 \tilde{\varphi}(z_2, \dots, z_n) &= (\ell_2 z_2, \dots, \ell_{t-1} z_{t-1}, \pm \mathbf{i} z_t, \ell_{t+1} z_{t+1}, \dots, \ell_{n-k} z_{n-k}, p^t(\mathbf{b} + DX)), \\
 \tilde{\varphi}^{-1}(z_2, \dots, z_n) &= (\bar{\ell}_2 z_2, \dots, \bar{\ell}_{t-1} z_{t-1}, \mp \mathbf{i} z_t, \bar{\ell}_{t+1} z_{t+1}, \dots, \bar{\ell}_{n-k} z_{n-k}, \\
 &\quad p^t(-D^{-1}\mathbf{b} + D^{-1}X)).
 \end{aligned}$$

Now, from (4.27), we consider

$$\begin{aligned}
 & \tilde{\varphi} g_2 \tilde{\varphi}^{-1}(z_2, \dots, z_n) \\
 &= (-z_2, \ell_3^2 \bar{z}_3, \dots, \ell_{t-1}^2 \bar{z}_{t-1}, -\bar{z}_t, \ell_{t+1}^2 \bar{z}_{t+1}, \dots, \ell_{n-k}^2 \bar{z}_{n-k}, \\
 &\quad \exp(4\pi \mathbf{i} b_{n-k+1}) \bar{z}_{n-k+1}, \dots, \exp(4\pi \mathbf{i} b_n) \bar{z}_n) \\
 &= g_2(z_2, \overline{\ell_3^2 z_3}, \dots, \overline{\ell_{t-1}^2 z_{t-1}}, -z_t, \overline{\ell_{t+1}^2 z_{t+1}}, \dots, \overline{\ell_{n-k}^2 z_{n-k}}, \\
 &\quad \exp(4\pi(-\mathbf{i})b_{n-k+1}) z_{n-k+1}, \dots, \exp(4\pi(-\mathbf{i})b_n) z_n) \\
 &= g_2 h(z_2, \dots, z_n)
 \end{aligned}$$

where

$$(4.41) \quad h(z_2, \dots, z_n) = (z_2, \overline{\ell_3^2 z_3}, \overline{\ell_4^2 z_4}, \dots, \overline{\ell_{t-1}^2 z_{t-1}}, -z_t, \overline{\ell_{t+1}^2 z_{t+1}}, \dots, \overline{\ell_{n-k}^2 z_{n-k}}, \\
 \exp(4\pi(-\mathbf{i})b_{n-k+1}) z_{n-k+1}, \dots, \exp(4\pi(-\mathbf{i})b_n) z_n).$$

We shall check that $h \notin \langle g_3, \dots, g_n \rangle$ (i.e., $g_2 h \notin (\mathbb{Z}_2)^{n-1}$).

Suppose that $h \in \langle g_3, \dots, g_n \rangle$. Since

$$h(z_{n-k+1}, \dots, z_n) = (\exp(4\pi(-\mathbf{i})b_{n-k+1}) z_{n-k+1}, \dots, \exp(4\pi(-\mathbf{i})b_n) z_n)$$

in (4.41), we may write $h = h'_{(even)}h''$ where $h' \in \langle g_3, \dots, g_{n-k} \rangle$ and $h'' \in \langle g_{n-k+1}, \dots, g_n \rangle$. Here $h'_{(even)}$ means a composition of an even number of generators $\{g_3, \dots, g_{n-k}\}$.

On the other hand, since $h(z_t) = -z_t$ in (4.41), we may write $h'_{(even)} = \hat{h}g_t\check{h}$, where $\hat{h}$ (resp. $\check{h}$) is a composition of an even (resp. odd) number of generators $\{g_3, \dots, g_{t-1}\}$ (resp. $\{g_{t+1}, \dots, g_{n-k}\}$). For $t = 4, 5, \dots, n-k-1$, such $h'_{(even)}$ implies that

$$h(z_{t+1}) = h'_{(even)}(z_{t+1}) = \begin{cases} \bar{z}_{t+1} & \text{if } h'_{(even)} = \hat{h}g_t\check{g}, \check{g} \in \langle g_{t+2}, \dots, g_{n-k} \rangle \\ -\bar{z}_{t+1} & \text{if } h'_{(even)} = \hat{h}g_tg_{t+1}\check{g}. \end{cases}$$

Hence this contradicts (4.41). Similarly for $h'_{(even)} = g_t\check{h}$ ($t = 4, 5, \dots, n-k-1$).

Now let us consider for $t = n-k$. Since $h(z_t) = -z_t$ in (4.41), $h = \dot{h}g_{n-k}h''$ where $\dot{h}$ is a composition of an even number of generators $\{g_3, \dots, g_{n-k-1}\}$. This implies that $h(z_{n-k+1}, \dots, z_n) = (\pm\bar{z}_{n-k+1}, \dots, \pm\bar{z}_n)$. This also contradicts (4.41). Similarly for $h = g_{n-k}h''$. Thus $g_2h \notin (\mathbb{Z}_2)^{n-1}$. Hence *Case 1* cannot occur.

Case 2. Let $g = g''$ where $g'' \in \langle g_{n-k+1}, \dots, g_n \rangle$.

If $g = g''$ satisfies (4.39), this implies that $(\widehat{z_4}^\alpha, \dots, \widehat{z_{n-k}}^\alpha) = (\widehat{z_4}^\beta, \dots, \widehat{z_{n-k}}^\beta)$. Then, from (4.38),

$$g''\beta(z_2, \dots, z_n) = (\bar{z}_2, z_3, \widehat{z_4}^\beta, \dots, \widehat{z_{n-k}}^\beta, \pm\bar{z}_{n-k+1}, \dots, \pm\bar{z}_{n-k+l-1}, \dots, \pm\bar{z}_{n-k+p-1}, \pm z_{n-k+p}, \dots, \pm z_n).$$

Then we obtain that the fixed point set of (α, T^{n-1}) is

$$Fix \alpha = (V, \{\pm 1\}_{n-k+1}, \dots, \{\pm 1\}_{n-k+l-1}, z_{n-k+l}, \dots, z_n)$$

with $V = \{(z_2, \dots, z_{n-k}) | \alpha(z_2, \dots, z_{n-k}) = (z_2, \dots, z_{n-k})\}$ and the fixed point set of $(g''\beta, T^{n-1})$ is

$$Fix g''\beta = (W, \{\star\}_{n-k+1}, \dots, \{\star\}_{n-k+p-1}, z_{n-k+p}, \dots, z_n)$$

with

$$W = \{(z_2, \dots, z_{n-k}) | g''\beta(z_2, \dots, z_{n-k}) = \beta(z_2, \dots, z_{n-k}) = (z_2, \dots, z_{n-k})\}$$

and $\star \in \{\pm 1, \pm \mathbf{i}\}$. Since $(\widehat{z_4}^\alpha, \dots, \widehat{z_{n-k}}^\alpha) = (\widehat{z_4}^\beta, \dots, \widehat{z_{n-k}}^\beta)$, $\dim V = \dim W$. Then by (4.26), we have

$$\begin{aligned} \dim(Fix \alpha) &= \dim(Fix g\beta) \\ \dim V + (k-l+1) &= \dim W + (k-p+1). \end{aligned}$$

Hence we get a contradiction. That is, the *Case 2* cannot occur. This completes the proof of Claim 1.

Claim 2. Bott matrices derived from each $A_{j'1}$ ($j = 1, \dots, k$) are not equivalent to each other, (i.e., there are $(2^{n-k-3} - 1)$ nonequivalent Bott matrices derived from each $A_{j'1}$ ($j = 1, \dots, k$)).

Associated with the entries $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k})$ of each $A_{j'1}$, there are $2^{(n-k-3)} - 1$ different actions $(\mathbb{Z}_2, M(B_1))_{j'}$. (Note that $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) \neq (0, \dots, 0)$.)

We prove that every two different actions $(\mathbb{Z}_2, M(B_1))_{j'}$ derived from $A_{j'1}$ (denoted by $(\alpha, M(B_1))_{j'_\alpha}$ and $(\beta, M(B_1))_{j'_\beta}$ respectively), the corresponding Bott matrices (denoted by $A_{j'_\alpha 1}$ and $A_{j'_\beta 1}$ respectively) are not equivalent.

Since the α -action and β -action are different, we may assume that $\widehat{z}_i^\alpha = z_i$, $\widehat{z}_i^\beta = \bar{z}_i$, for some $i \in \{4, \dots, n-k\}$. Then

$$(4.42) \quad \begin{aligned} & \alpha(z_2, \dots, z_n) = \\ & (\bar{z}_2, z_3, \widehat{z}_4^\alpha, \dots, \widehat{z}_{i-1}^\alpha, z_i, \widehat{z}_{i+1}^\alpha, \dots, \widehat{z}_{n-k}^\alpha, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+j-1}, z_{n-k+j}, \dots, z_n), \\ & \beta(z_2, \dots, z_n) = \\ & (\bar{z}_2, z_3, \widehat{z}_4^\beta, \dots, \widehat{z}_{i-1}^\beta, \bar{z}_i, \widehat{z}_{i+1}^\beta, \dots, \widehat{z}_{n-k}^\beta, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+j-1}, z_{n-k+j}, \dots, z_n), \\ & \text{for } i = 4, \dots, n-k, j = 1, \dots, k. \end{aligned}$$

As before, if $A_{j'_\alpha 1}$ and $A_{j'_\beta 1}$ are equivalent, we have

$$(4.43) \quad \begin{aligned} & \tilde{\varphi}(\alpha(z_2, \dots, z_n)) = g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ & (\ell_2\bar{z}_2, \ell_3z_3, \ell_4\widehat{z}_4^\alpha, \dots, \ell_{i-1}\widehat{z}_{i-1}^\alpha, \ell_i z_i, \ell_{i+1}\widehat{z}_{i+1}^\alpha, \dots, \ell_{n-k}\widehat{z}_{n-k}^\alpha, c_{n-k+1}v'_{n-k+1}, \\ & \dots, c_n v'_n) \\ & = g(\overline{\ell_2\bar{z}_2}, \overline{\ell_3z_3}, \overline{\ell_4\widehat{z}_4^\beta}, \dots, \overline{\ell_{i-1}\widehat{z}_{i-1}^\beta}, \overline{\ell_i z_i}, \overline{\ell_{i+1}\widehat{z}_{i+1}^\beta}, \dots, \overline{\ell_{n-k}\widehat{z}_{n-k}^\beta}, \\ & \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j-1}w_{n-k+j-1}}, c_{n-k+j}w_{n-k+j}, \dots, c_n w_n), \end{aligned}$$

for some $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$, where

$$(4.44) \quad (v'_{n-k+1}, \dots, v'_n) = p((-x_{n-k+1}, \dots, -x_{n-k+j-1}, x_{n-k+j}, \dots, x_n)^t D).$$

Obviously, $g \in \langle g_2, g_3 \rangle$ does not satisfy (4.43) because of (4.40). If $g \in \langle g_i, \dots, g_n \rangle$ then the equation (4.43) is also not satisfied, because it implies that

$$(4.45) \quad \ell_i z_i = g(\overline{\ell_i z_i}) = \begin{cases} -\overline{\ell_i z_i} & \text{if } g = g_i g' \text{ where } g' = \langle g_{i+1}, \dots, g_n \rangle \\ \overline{\ell_i z_i} & \text{if } g = g'. \end{cases}$$

Because of (4.40) and (4.45), if $i = 4$ then there is no $g \in (\mathbb{Z}_2)^{n-1}$ satisfying (4.43).

Now let us consider for $i \in \{5, \dots, n-k\}$. Since $\widehat{z}_i^\alpha = z_i$ and $\widehat{z}_i^\beta = \bar{z}_i$, for some $i \in \{5, \dots, n-k\}$, we may write $g = g_t \dot{g}^{(even)} \ddot{g}$ with $g_t \in \{g_4, \dots, g_{i-1}\}$ ($t = 4, \dots, i-1$), $\dot{g} \in \langle g_{t+1}, \dots, g_{i-1} \rangle$, $\ddot{g} \in \langle g_{i+1}, \dots, g_n \rangle$ (if $t = i-1$ then $g = g_t \ddot{g}$). Here $\dot{g}^{(even)}$ means a composition of an even number of generators of $\dot{g}$.

Note that since $g_t(z_t) = -z_t$, $g(z_t) = -z_t$. If $\widehat{z}_t^\alpha \neq \widehat{z}_t^\beta$ (resp. $\widehat{z}_t^\alpha = \widehat{z}_t^\beta = z_t$) then $g = g_t \dot{g}^{(even)} \ddot{g}$ does not satisfy (4.43), because it implies that $\ell_t z_t = g(\overline{\ell_t z_t}) = -\overline{\ell_t z_t}$ (resp. $\ell_t z_t = g(\ell_t z_t) = -\ell_t z_t$). If $\widehat{z}_t^\alpha = \widehat{z}_t^\beta = \bar{z}_t$ then $\ell_t \bar{z}_t = g(\overline{\ell_t \bar{z}_t}) = -\overline{\ell_t \bar{z}_t}$. This implies that $\ell_t = \pm \mathbf{i}$. Therefore

$$\tilde{\varphi}(z_2, \dots, z_n) = (\ell_2 z_2, \dots, \ell_{t-1} z_{t-1}, \pm \mathbf{i} z_t, \ell_{t+1} z_{t+1}, \dots, \ell_{n-k} z_{n-k}, p^t(\mathbf{b} + DX)).$$

Similar to the proof of Claim 1, one can check that $\tilde{\varphi} g_2 \tilde{\varphi}^{-1}(z_2, \dots, z_n) \notin (\mathbb{Z}_2)^{n-1}$. Hence we have a contradiction. That is, there is no $g \in (\mathbb{Z}_2)^{n-1}$ satisfying (4.43).

Since all combinations of $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k})$ with $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) \neq (0, \dots, 0)$ are different for each $A_{j'1}$, there are $(2^{n-k-3} - 1)$ nonequivalent Bott matrices derived from each $A_{j'1}$ ($j = 1, \dots, k$).

Claim 3. Bott matrices derived from $A_{(k+1)'1}$ are not equivalent to each other, (i.e., there are $2^{n-k-3} - (n - k - 2)$ nonequivalent Bott matrices derived from $A_{(k+1)'1}$).

Since there are $2^{n-k-3} - (n - k - 2)$ different combination of $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k})$ with

$$(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) \neq \overbrace{(0, \dots, 0, 1, \dots, 1)}^l,$$

(that is, there are $2^{n-k-3} - (n - k - 2)$ different actions $(\mathbb{Z}_2, M(B_1))_{(k+1)'}$), by using argument in the proof of Claim 2 above, there are $2^{n-k-3} - (n - k - 2)$ nonequivalent Bott matrices derived from $A_{(k+1)'1}$.

According to Claim 1, 2, 3, we obtain that there are $k(2^{n-k-3} - 1) + (2^{n-k-3} - (n - k - 2))$ nonequivalent Bott matrices derived from (4.36). $\square$

Remark 4.10. Consider Bott matrices A_{i1} ($i = 1, \dots, n - 2$) in (4.22) and $A_{j'1}$ ($j = 1, \dots, k, k + 1$) in (4.36).

- (i) Associated with the entries $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k})$ in $A_{j'1}$ ($j = 1, \dots, k$), if $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) = (0, \dots, 0)$ then $A_{1'1} = A_{11}$ and $A_{j'1} \sim A_{(n-j)1}$ ($j = 2, \dots, k$) by the equivariant diffeomorphism

$$(\Phi, \varphi): (\mathbb{Z}_2, M(B_1))_{j'} \rightarrow (\mathbb{Z}_2, M(B_1))_{n-j}$$

defined by

$$\begin{aligned} [z_2, \dots, z_{n-k}, z_{n-k+1}, \dots, z_{n-k+(j-1)}, z_{n-k+j}, \dots, z_n] &\xrightarrow{\varphi} \\ [\mathbf{i}z_2, \dots, z_{n-k}, z_{n-k+j}, \dots, z_n, z_{n-k+1}, \dots, z_{n-k+(j-1)}]. \end{aligned}$$

- (ii) Associated with the entries $(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k})$ in $A_{(k+1)'1}$, if

$$(\{\hat{1}\}_4, \dots, \{\hat{1}\}_{n-k}) = \overbrace{(0, \dots, 0, 1, \dots, 1)}^l \quad (l = 0, 1, \dots, n-k-3), \text{ then } A_{(k+1)'1} \sim A_{(l+2)1} \quad (l = 0, 1, \dots, n-k-3) \text{ by the equivariant diffeomorphism}$$

$$(\Phi, \varphi): (\mathbb{Z}_2, M(B_1))_{(k+1)'} \rightarrow (\mathbb{Z}_2, M(B_1))_{(l+2)}$$

defined by

$$\varphi([z_2, \dots, z_n]) = [\mathbf{i}z_2, \dots, z_n].$$

Lemma 4.11. Each Bott matrix in (4.22) is not equivalent to each Bott matrix derived from (4.36).

Proof. Using the argument at the beginning of the proof of Lemma (4.8) we shall prove the following Claims.

Claim i). A_{l1} ($l = 1, \dots, n - k - 1$) is not equivalent to $A_{1'1}$.

For this, we prove the following cases.

Case 1. A_{11} is not equivalent to $A_{1'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_1$ and $(\mathbb{Z}_2, M(B_1))_{1'}$ are

$$(4.46) \quad \alpha[z_2, \dots, z_n] = [\bar{z}_2, z_3, \dots, z_n]$$

and

$$(4.47) \quad \beta[z_2, \dots, z_n] = [\bar{z}_2, z_3, \hat{z}_4, \dots, \hat{z}_{n-k}, z_{n-k+1}, \dots, z_n]$$

respectively. If A_{11} is equivalent to $A_{1'1}$, by (4.26), there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.48) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \ell_3z_3, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n). \end{aligned}$$

As before, $g \in \langle g_2, g_3 \rangle$ does not satisfy (4.48) because of (4.40). Since

$$(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) = g(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n),$$

g is a composition of an even number of generators $\{g_4, \dots, g_{n-k}\}$ and $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not satisfy (4.48). Then we can take $g = g_t g'$ with $t = 4, \dots, n-k-1$ and $g' \in \langle g_{t+1}, \dots, g_{n-k} \rangle$. This implies that

$$\ell_t z_t = g(\widehat{\ell_t z_t}) = \begin{cases} -\ell_t z_t & \text{if } \widehat{\ell_t z_t} = \ell_t z_t \\ -\overline{\ell_t z_t} & \text{if } \widehat{\ell_t z_t} = \overline{\ell_t z_t}. \end{cases}$$

Therefore there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.48).

Case 2. A_{j1} ($j = 2, \dots, n-k-1$) is not equivalent to $A_{1'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_j$ and $(\mathbb{Z}_2, M(B_1))_{1'}$ are

$$(4.49) \quad \begin{aligned} \alpha[z_2, \dots, z_n] &= [\bar{z}_2, \bar{z}_3, \dots, \bar{z}_{j+1}, z_{j+2}, \dots, z_n] \\ &= [g_2 g_{j+1}(\bar{z}_2, \bar{z}_3, \dots, \bar{z}_{j+1}, z_{j+2}, \dots, z_n)] \\ &= [-\bar{z}_2, z_3, \dots, z_j, -z_{j+1}, z_{j+2}, \dots, z_n] \end{aligned}$$

and as in (4.47) respectively. If A_{j1} is equivalent to $A_{1'1}$, there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.50) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (-\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_j z_j, -\ell_{j+1}z_{j+1}, \ell_{j+2}z_{j+2}, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \\ &\dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \ell_3z_3, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n). \end{aligned}$$

Similar to the argument of the proof of *Case 1* above, $g \in \langle g_2, \dots, g_j \rangle$ ($j = 2, \dots, n-k-1$) does not satisfy (4.50). Since

$$(4.51) \quad (c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) = g(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n)$$

in (4.50), $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not satisfy (4.50). Because of

$$(-\ell_{j+1}z_{j+1}) = \begin{cases} g(\ell_{j+1}z_{j+1}) & \text{if } j = 2 \\ g(\widehat{\ell_{j+1}z_{j+1}}) & \text{if } j = 3, \dots, n-k-1 \end{cases}$$

and (4.51), then g is a composition of an even number of generators $\{g_{j+1}, \dots, g_{n-k}\}$ which can be written by $g = g_{j+1}\dot{g}$ with $j = 2, \dots, n-k-2$, $\dot{g} \in \langle g_{j+2}, \dots, g_{n-k} \rangle$. However this implies that

$$\ell_t z_t = g(\widehat{\ell_t z_t}) = -\widehat{\ell_t z_t}$$

for some t ($j+2 \leq t \leq n-k$). For $j = n-k-1$, $g = g_{n-k}$ does not satisfy (4.50), because by assumption (4.37), $(\hat{z}_4, \dots, \hat{z}_n) \neq (z_4, \dots, z_n)$ in (4.47). Hence there is

no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.50). This completes the proof of Claim i).

Claim ii). A_{l1} ($l = 1, \dots, n - k - 1$) is not equivalent to $A_{j'1}$ ($j = 2, \dots, k$).

For this, we prove the following cases.

Case 1. A_{11} is not equivalent to $A_{j'1}$ ($j = 2, \dots, k$).

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_1$ and $(\mathbb{Z}_2, M(B_1))_{j'}$ are as in (4.46) and

$$(4.52) \quad \beta[z_2, \dots, z_n] = [\bar{z}_2, z_3, \hat{z}_4, \dots, \hat{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{n-k+j-1}, z_{n-k+j}, \dots, z_n]$$

respectively. If A_{11} is equivalent to $A_{j'1}$, there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.53) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \overline{\ell_3z_3}, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j-1}w_{n-k+j-1}}, \\ &\quad c_{n-k+j}w_{n-k+j}, \dots, c_nw_n). \end{aligned}$$

Since

$$(4.54) \quad \begin{aligned} (c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) &= \\ g(\overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j-1}w_{n-k+j-1}}, c_{n-k+j}w_{n-k+j}, \dots, c_nw_n), \end{aligned}$$

it is clear that there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.53).

Case 2. A_{i1} ($i = 2, \dots, n - k - 1$) is not equivalent to $A_{j'1}$ ($j = 2, \dots, k$).

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_i$ and $(\mathbb{Z}_2, M(B_1))_{j'}$ are as in (4.49) and (4.52) respectively. If A_{i1} is equivalent to $A_{j'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.55) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (-\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_i z_i, -\ell_{i+1}z_{i+1}, \ell_{i+2}z_{i+2}, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, \\ c_nw_n) \\ &= g(\overline{\ell_2z_2}, \overline{\ell_3z_3}, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_{n-k+j-1}w_{n-k+j-1}}, \\ &\quad c_{n-k+j}w_{n-k+j}, \dots, c_nw_n). \end{aligned}$$

Because of (4.54), there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.55). This completes the proof of Claim ii).

Claim iii). A_{l1} ($l = 1, \dots, n - k - 1$) is not equivalent to $A_{(k+1)'1}$.

For this, we prove the following cases.

Case 1. A_{11} is not equivalent to $A_{(k+1)'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_1$ and $(\mathbb{Z}_2, M(B_1))_{(k+1)'}$ are as in (4.46) and

$$(4.56) \quad \beta[z_2, \dots, z_n] = [\bar{z}_2, z_3, \hat{z}_4, \dots, \hat{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_n]$$

respectively. If A_{11} is equivalent to $A_{(k+1)'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.57) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \\ &= g(\overline{\ell_2z_2}, \overline{\ell_3z_3}, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n}). \end{aligned}$$

As before, $g \in \langle g_2, g_3 \rangle$ does not satisfy (4.57) because of (4.40). Since

$$(c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) = g(\overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n}),$$

g is a composition of an odd number of generators $\{g_4, \dots, g_{n-k}\}$ and $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not satisfy (4.57). Then we can take $g = g_t g'$, with $t = 4, \dots, n-k-1$ and $g' \in \langle g_{t+1}, \dots, g_{n-k} \rangle$. This implies that

$$\ell_t z_t = g(\widehat{\ell_t z_t}) = \begin{cases} -\ell_t z_t & \text{if } \widehat{\ell_t z_t} = \ell_t z_t \\ -\overline{\ell_t z_t} & \text{if } \widehat{\ell_t z_t} = \overline{\ell_t z_t}. \end{cases}$$

For $t = n-k$, it is clear that $g = g_{n-k}$ does not also satisfy (4.57). Hence there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.57).

Case 2. A_{j1} ($j = 2, \dots, n-k-2$) is not equivalent to $A_{(k+1)'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_j$ and $(\mathbb{Z}_2, M(B_1))_{(k+1)'}$ are as in (4.49) and (4.56) respectively. If A_{j1} is equivalent to $A_{(k+1)'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.58) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (-\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_j z_j, -\ell_{j+1}z_{j+1}, \ell_{j+2}z_{j+2}, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}w_{n-k+1}, \dots, \\ c_nw_n) \\ &= g(\overline{\ell_2z_2}, \overline{\ell_3z_3}, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n}). \end{aligned}$$

Note that by assumption (see (4.37)),

$$(4.59) \quad \begin{aligned} (z_3, \hat{z}_4, \dots, \hat{z}_{j+1}, \hat{z}_{j+2}, \dots, \hat{z}_{n-k}) &\neq (z_3, z_4, \dots, z_{j+1}, \bar{z}_{j+2}, \dots, \bar{z}_{n-k}), \\ (\hat{z}_4, \dots, \hat{z}_{n-k}) &\neq (z_4, \dots, z_{n-k}) \end{aligned}$$

in (4.56) for $j = 2, \dots, n-k-2$.

Similar to the argument of the proof of *Case 1* above, $g \in \langle g_2, \dots, g_j \rangle$ ($j = 2, \dots, n-k-1$) does not satisfy (4.58). Since

$$(4.60) \quad (c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) = g(\overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n})$$

in (4.58), $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not satisfy (4.58). Because of

$$(-\ell_{j+1}z_{j+1}) = \begin{cases} g(\ell_{j+1}z_{j+1}) & \text{if } j = 2 \\ g(\widehat{\ell_{j+1}z_{j+1}}) & \text{if } j = 3, \dots, n-k-2 \end{cases}$$

and (4.60), then g is a composition of an odd number of generators $\{g_{j+1}, \dots, g_{n-k}\}$ which can be written as $g = g_{j+1}\dot{g}$ ($j = 2, \dots, n-k-2$), $\dot{g} \in \langle g_{j+2}, \dots, g_{n-k} \rangle$. However this implies that

$$\ell_t z_t = g(\widehat{\ell_t z_t}) = -\widehat{\ell_t z_t}$$

for some t ($j+2 \leq t \leq n-k$). Hence such g does not satisfy (4.58). In particular, $g = g_{j+1}$ ($j = 2, \dots, n-k-1$) does not also satisfy (4.58) because of (4.59). Thus

there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.58).

Case 3. $A_{(n-k-1)1}$ is not equivalent to $A_{(k+1)'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_{(n-k-1)}$ and $(\mathbb{Z}_2, M(B_1))_{(k+1)'}$ are

$$\begin{aligned} \alpha[z_2, \dots, z_n] &= [\bar{z}_2, \bar{z}_3, \dots, \bar{z}_{n-k}, z_{n-k+1}, \dots, z_n] \\ (4.61) \quad &= [g_2(\bar{z}_2, \bar{z}_3, \dots, \bar{z}_{n-k}, z_{n-k+1}, \dots, z_n)] \\ &= [-\bar{z}_2, z_3, \dots, z_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_n] \end{aligned}$$

and as in (4.56) respectively. If $A_{(n-k-1)1}$ is equivalent to $A_{(k+1)'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$\begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (4.62) \quad &(-\ell_2\bar{z}_2, \ell_3z_3, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}\bar{w}_{n-k+1}, \dots, c_n\bar{w}_n) \\ &= g(\ell_2\bar{z}_2, \ell_3z_3, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, \overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n}). \end{aligned}$$

Note that by assumption (4.37), $(\hat{z}_4, \dots, \hat{z}_{n-k}) \neq (z_4, \dots, z_{n-k})$ in (4.56). Therefore $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not satisfy (4.62).

Since

$$(c_{n-k+1}\bar{w}_{n-k+1}, \dots, c_n\bar{w}_n) = g(\overline{c_{n-k+1}w_{n-k+1}}, \dots, \overline{c_nw_n}),$$

g is a composition of an even number of generators $\{g_4, \dots, g_{n-k}\}$. Similar to the argument in *Case 1* (Claim i)), such g does not satisfy (4.62). Thus there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.62). This completes the proof of Claim iii).

Claim iv). A_{l1} ($l = n-k, \dots, n-2$) is not equivalent to $A_{j'1}$ ($j = 1, \dots, k, k+1$).

For this, we prove the following cases.

Case 1. A_{l1} ($l = n-k, \dots, n-2$) is not equivalent to $A_{1'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_l$ and $(\mathbb{Z}_2, M(B_1))_{1'}$ are

$$\begin{aligned} \alpha[z_2, \dots, z_n] &= [\bar{z}_2, \bar{z}_3, \bar{z}_4, \dots, \bar{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{1+l}, z_{2+l}, \dots, z_n] \\ (4.63) \quad &= [g_2(\bar{z}_2, \bar{z}_3, \bar{z}_4, \dots, \bar{z}_{n-k}, \bar{z}_{n-k+1}, \dots, \bar{z}_{1+l}, z_{2+l}, \dots, z_n)] \\ &= [-\bar{z}_2, z_3, z_4, \dots, z_{n-k}, z_{n-k+1}, \dots, z_{1+l}, \bar{z}_{2+l}, \dots, \bar{z}_n] \end{aligned}$$

and as in (4.47) respectively. If A_{l1} is equivalent to $A_{1'1}$, there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$\begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (4.64) \quad &(-\ell_2\bar{z}_2, \ell_3z_3, \ell_4z_4, \dots, \ell_{n-k}z_{n-k}, c_{n-k+1}y'_{n-k+1}, \dots, c_ny'_n) \\ &= g(\ell_2\bar{z}_2, \ell_3z_3, \widehat{\ell_4z_4}, \dots, \widehat{\ell_{n-k}z_{n-k}}, c_{n-k+1}w_{n-k+1}, \dots, c_nw_n) \end{aligned}$$

where

$$(y'_{n-k+1}, \dots, y'_n) = p((x_{n-k+1}, \dots, x_{1+l}, -x_{2+l}, \dots, -x_n)^t D).$$

Note that by assumption (4.37),

$$(\hat{z}_4, \dots, \hat{z}_{n-k}) \neq (z_4, \dots, z_{n-k})$$

in (4.47). Therefore $g \in \langle g_{n-k+1}, \dots, g_n \rangle$ does not occur. As before $g \in \langle g_2, g_3 \rangle$ does not also satisfy (4.64).

Now we consider $g \in \langle g_4, \dots, g_n \rangle$. We may write $g = g_t g' g''$ with $t = 4, \dots, n - k - 1$, $g' \in \langle g_{t+1}, \dots, g_{n-k} \rangle$, $g'' \in \langle g_{n-k+1}, \dots, g_n \rangle$ (if $t = n - k$ then $g = g_{n-k} g''$). This implies that

$$\ell_t z_t = g(\widehat{\ell_t z_t}) = \begin{cases} -\ell_t z_t & \text{if } \widehat{\ell_t z_t} = \ell_t z_t \\ -\overline{\ell_t z_t} & \text{if } \widehat{\ell_t z_t} = \overline{\ell_t z_t}. \end{cases}$$

Therefore there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.64).

Case 2. A_{l1} ($l = n - k, \dots, n - 2$) is not equivalent to $A_{j'1}$ ($j = 2, \dots, k$).

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_l$ and $(\mathbb{Z}_2, M(B_1))_{j'}$ are as in (4.63) and (4.52) respectively. If A_{l1} is equivalent to $A_{j'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.65) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (-\ell_2 \bar{z}_2, \ell_3 z_3, \ell_4 z_4, \dots, \ell_{n-k} z_{n-k}, c_{n-k+1} y'_{n-k+1}, \dots, c_n y'_n) \\ &= g(\ell_2 z_2, \ell_3 z_3, \widehat{\ell_4 z_4}, \dots, \widehat{\ell_{n-k} z_{n-k}}, \overline{c_{n-k+1} w_{n-k+1}}, \dots, \overline{c_{n-k+j-1} w_{n-k+j-1}}, \\ &\quad c_{n-k+j} w_{n-k+j}, \dots, c_n w_n). \end{aligned}$$

Note that

$$(\hat{z}_4, \dots, \hat{z}_{n-k}) \neq (z_4, \dots, z_{n-k})$$

in (4.52). Similar to the previous argument, there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.65).

Case 3. A_{l1} ($l = n - k, \dots, n - 2$) is not equivalent to $A_{(k+1)'1}$.

By the definition, the actions $(\mathbb{Z}_2, M(B_1))_l$ and $(\mathbb{Z}_2, M(B_1))_{(k+1)'}$ are as in (4.63) and (4.56) respectively. If A_{l1} is equivalent to $A_{(k+1)'1}$ then there is $g \in (\mathbb{Z}_2)^{n-1}$ such that

$$(4.66) \quad \begin{aligned} \tilde{\varphi}(\alpha(z_2, \dots, z_n)) &= g\beta\tilde{\varphi}(z_2, \dots, z_n), \\ (-\ell_2 \bar{z}_2, \ell_3 z_3, \ell_4 z_4, \dots, \ell_{n-k} z_{n-k}, c_{n-k+1} y'_{n-k+1}, \dots, c_n y'_n) \\ &= g(\ell_2 z_2, \ell_3 z_3, \widehat{\ell_4 z_4}, \dots, \widehat{\ell_{n-k} z_{n-k}}, \overline{c_{n-k+1} w_{n-k+1}}, \dots, \overline{c_n w_n}). \end{aligned}$$

Note that by assumption

$$(\hat{z}_4, \dots, \hat{z}_{n-k}) \neq (z_4, \dots, z_{n-k})$$

in (4.56). Similar to the argument in *Case 1* (Claim iv)) above, there is no $g \in (\mathbb{Z}_2)^{n-1} = \langle g_2, \dots, g_n \rangle$ satisfying (4.66). This completes the proof of Claim iv). $\square$

Theorem 4.12. *Let $M(A) = S^1 \times_{\mathbb{Z}_2} M(B)$ be an n -dimensional real Bott manifold. Suppose that B is either one of the list in (4.67). Then $M(B)$ are diffeomorphic each other and the number of diffeomorphism classes of such real Bott manifolds $M(A)$ above is $(k+1)2^{n-k-3}$ ($k \geq 2$, $n-k \geq 3$).*

$$\begin{aligned}
(4.67) \quad & B_1 = \begin{pmatrix} 1 & 1 & 1 & \dots & \dots & \dots & 1 \\ & 1 & 1 & \dots & \dots & \dots & 1 \\ & & \ddots & & & & \vdots \\ & & & 1 & 1 & \dots & 1 \\ 0 & & & & & I_k & \end{pmatrix}, B_2 = \begin{pmatrix} 1 & 1 & 0 & \dots & \dots & \dots & 0 \\ & 1 & 1 & \dots & \dots & \dots & 1 \\ & & \ddots & & & & \vdots \\ & & & 1 & 1 & \dots & 1 \\ 0 & & & & & I_k & \end{pmatrix}, \\
& \vdots \\
& B_{n-k-1} = \begin{pmatrix} 1 & 1 & 0 & \dots & \dots & \dots & \dots & 0 \\ & 1 & 1 & 0 & \dots & \dots & \dots & 0 \\ & & \ddots & & & & & \vdots \\ & & & 1 & 1 & 0 & \dots & 0 \\ & & & & 1 & 1 & \dots & 1 \\ 0 & & & & & & I_k & \end{pmatrix}, \\
& B_{n-k} = \begin{pmatrix} 1 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 \\ & 1 & 1 & 0 & \dots & \dots & \dots & \dots & 0 \\ & & \ddots & & & & & & \vdots \\ & & & 1 & 1 & 0 & 0 & 0 & \dots & 0 \\ & & & & 1 & 1 & 1 & 0 & \dots & 0 \\ & & & & & 1 & 1 & 0 & \dots & 0 \\ & & & & & & 1 & 1 & \dots & 1 \\ 0 & & & & & & & & I_k & \end{pmatrix}, \\
& \vdots \\
& B_{n-k+(n-k-4)} = \begin{pmatrix} 1 & 1 & \dots & \dots & 1 & 0 & \dots & 0 \\ & 1 & \dots & \dots & 1 & 0 & \dots & 0 \\ & & \ddots & & \vdots & \vdots & & \vdots \\ & & & 1 & 1 & 0 & \dots & 0 \\ & & & & 1 & 1 & \dots & 1 \\ 0 & & & & & & I_k & \end{pmatrix}.
\end{aligned}$$

Proof. Note that each B_j is of size $n-1$ and $M(B_j) = S^1 \times_{\mathbb{Z}_2} M(B'_j)$ for some $(n-2)$ -dimensional real Bott manifold $M(B'_j)$. $M(B_{j-1})$ is diffeomorphic to $M(B_j)$ ($j = 2, 3, \dots, n-k-1$) by the equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B'_{j-1})) \rightarrow (\mathbb{Z}_2, M(B'_j))$ which is defined by $\varphi([z_2, \dots, z_{n-1}]) = [z_2, \dots, \mathbf{i}z_j, \dots, z_{n-1}]$. $M(B_a)$ is diffeomorphic to $M(B_b)$ ($a = n-k + (n-k-3) - (j-1)$; $b = n-k + (n-$

$k-3)-(j-2)$; $j = n-k-1, n-k-2, \dots, 4, 3$) by the equivariant diffeomorphism $\varphi : (\mathbb{Z}_2, M(B'_a)) \rightarrow (\mathbb{Z}_2, M(B'_b))$ which is defined by $\varphi([z_2, \dots, z_{n-1}]) = [z_2, \dots, \mathbf{i}z_{j-1}, \dots, z_{n-1}]$.

By the hypothesis, there are 2^{n-2} possible $\mathbb{Z}_2$ -actions on each $M(B_j)$, $j = 1, \dots, 2(n-k)-4$. We shall prove that among $2^{n-2} \times (2(n-k)-4)$ real Bott manifolds created from $M(B_j)$ ($j = 1, \dots, 2(n-k)-4$), there are only $(k+1)2^{n-k+3}$ diffeomorphism classes.

First of all we show that there are $(k+1)2^{n-k+3}$ diffeomorphism classes of real Bott manifolds $M(A_{i1})$ created from $M(B_1)$ by $\mathbb{Z}_2$ -actions.

If a Bott matrix A' created from B_1 is different from the Bott matrices in (4.22) and (4.36), it is easy to check that the corresponding $(\mathbb{Z}_2, M(B_1))$ is equivariantly diffeomorphic to one of the actions $(\mathbb{Z}_2, M(B_1))$ corresponding to the Bott matrices in (4.22) or (4.36) by the ad hoc argument. (Compare Section 2 for the argument to find an equivariant diffeomorphism.) Once there exists such an equivariant diffeomorphism, A' is equivalent to one of A_{i1} 's ($i = 1, \dots, n-2$) in (4.22) or $A_{j'1}$'s ($j = 1, \dots, k, (k+1)$) in (4.36) by Theorem 1.4. Then, because of Lemma 4.8, 4.9, 4.11 and Remark 4.10, there are $(n-2) + (2^{n-k-3} - 1)k + 2^{n-k-3} - (n-k-2) = (k+1)2^{n-k+3}$ distinct diffeomorphism classes of $M(A_{i1})$ ($i = 1, \dots, 2^{n-2}$) created from $M(B_1)$.

Next, we show that $M(A_{ij})$ ($i = 1, \dots, 2^{n-2}$) created from $M(B_j)$ ($j = 2, \dots, n-k + (n-k-4)$) is diffeomorphic to one of the real Bott manifolds corresponding to Bott matrices in (4.22) or (4.36).

For brevity we can consider

$$(4.68) \quad A_{\ell 1} = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(1)}} & \dots & \dots & \dots & \{\hat{1}\}_{n_{(1)}} \\ \hline 0 & & & & & & & B_1 \end{array} \right)$$

($\ell = 1, \dots, n-2, 1', \dots, k', (k+1)'$) representing each $A_{j'1}$ ($j = 1, \dots, k, (k+1)$) in (4.36), and each A_{i1} ($i = 1, \dots, n-2$) in (4.22), where A_{i1} ($i = 2, \dots, n-2$) is equivalent to (4.68) by the equivariant diffeomorphism $\varphi : (\mathbb{Z}_2, M(B_1)) \rightarrow (\mathbb{Z}_2, M(B_1))$ which is defined by $\varphi([z_2, \dots, z_n]) = [\mathbf{i}z_2, \dots, z_n]$. Note that $\{\hat{1}\}_{l_{(j)}}$ means $\hat{1} \in \{0, 1\}$ in the l -th spot where the corresponding Bott matrix is created from B_j .

Now we define an equivariant diffeomorphism $\varphi : (\mathbb{Z}_2, M(B_1)) \rightarrow (\mathbb{Z}_2, M(B_2))$ by $\varphi([z_2, \dots, z_n]) = [z_2, \mathbf{i}z_3, z_4, \dots, z_n]$, then (4.68) is equivalent to

$$(4.69) \quad A_{\ell 2} = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(2)}} & \dots & \dots & \dots & \{\hat{1}\}_{n_{(2)}} \\ \hline 0 & & & & & & & B_2 \end{array} \right).$$

Next we define an equivariant diffeomorphism $\varphi : (\mathbb{Z}_2, M(B_2)) \rightarrow (\mathbb{Z}_2, M(B_3))$ by $\varphi([z_2, \dots, z_n]) = [z_2, z_3, \mathbf{i}z_4, z_5, \dots, z_n]$ so that Bott matrix (4.69) is equivalent to

$$A_{\ell 3} = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(3)}} & \dots & \dots & \dots & \{\hat{1}\}_{n_{(3)}} \\ \hline 0 & & & & & & & B_3 \end{array} \right).$$

In general, let us consider

(4.70)

$$A_{\ell(j-1)} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(j-1)}} & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \{\hat{1}\}_{n_{(j-1)}} & \\ \hline & 1 & 1 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & \\ & & 1 & 1 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & \\ & & & \ddots & & & & & & & \vdots & \\ & & & & 1 & 1 & 0 & \cdots & \cdots & \cdots & 0 & \\ & 0 & & & & 1 & 1 & \cdots & \cdots & \cdots & 1 & \\ & & & & & & 1 & 1 & \cdots & \cdots & 1 & \\ & & & 0 & & & \ddots & & & & \vdots & \\ & & & & & & & 1 & 1 & \cdots & 1 & \\ & & & & & & & & I_k & & & \end{array} \right) \begin{array}{l} g_1 \\ g_2 \\ g_3 \\ \vdots \\ g_{j-1} \\ g_j \\ g_{j+1} \\ \vdots \\ g_{n-k} \end{array}$$

($j = 2, \dots, n-k-1$). Defining an equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B_{j-1})) \rightarrow (\mathbb{Z}_2, M(B_j))$ by $\varphi([z_2, \dots, z_n]) = [z_2, \dots, z_j, \mathbf{i}z_{j+1}, z_{j+2}, \dots, z_n]$, we obtain that (4.70) is equivalent to

$$A_{\ell(j)} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(j)}} & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \{\hat{1}\}_{n_{(j)}} & \\ \hline & 1 & 1 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & \\ & & 1 & 1 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & \\ & & & \ddots & & & & & & & \vdots & \\ & & & & 1 & 1 & 0 & \cdots & \cdots & \cdots & 0 & \\ & 0 & & & & 1 & 1 & 0 & \cdots & \cdots & 0 & \\ & & & & & & 1 & 1 & \cdots & \cdots & 1 & \\ & & & 0 & & & \ddots & & & & \vdots & \\ & & & & & & & 1 & 1 & \cdots & 1 & \\ & & & & & & & & I_k & & & \end{array} \right) \begin{array}{l} g_1 \\ g_2 \\ g_3 \\ \vdots \\ g_{j-1} \\ g_j \\ g_{j+1} \\ \vdots \\ g_{n-k} \end{array}.$$

Therefore the previous Bott matrix is equivalent to

(4.71)

$$A_{\ell(n-k-1)} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(n-k-1)}} & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \{\hat{1}\}_{n_{(n-k-1)}} & \\ \hline 0 & & & & B_{n-k-1} & & & & & & & \end{array} \right).$$

Next, (4.71) is equivalent to the following one by the equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B_{n-k-1})) \rightarrow (\mathbb{Z}_2, M(B_{n-k}))$ defined by $\varphi([z_2, \dots, z_n]) = [z_2, \dots, z_{n-k-2}, \mathbf{i}z_{n-k-1}, z_{n-k}, \dots, z_n]$

$$A_{\ell(n-k)} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4_{(n-k)}} & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \{\hat{1}\}_{n_{(n-k)}} & \\ \hline 0 & & & & B_{n-k} & & & & & & & \end{array} \right).$$

In general, let us consider the following Bott matrix

(4.72)

$$A_{\ell a} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4(a)} & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \{\hat{1}\}_{n(a)} \\ \hline & 1 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ & & 1 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ & & & \ddots & & & & & & & & \vdots \\ & & & & 1 & 1 & 0 & \dots & 0 & 0 & \dots & 0 \\ & & & & & 1 & 1 & \dots & 1 & 0 & \dots & 0 \\ & & & & & & 1 & \dots & 1 & 0 & \dots & 0 \\ & & & & & & \ddots & & & & & \vdots \\ & & & & & & & 1 & 1 & 0 & \dots & 0 \\ & & & & & & & & 1 & 1 & \dots & 1 \\ & & & & & & & & & & & I_k \end{array} \right) \begin{array}{l} g_1 \\ g_2 \\ g_3 \\ \vdots \\ g_{j-1} \\ g_j \\ g_{j+1} \\ \vdots \\ g_{n-k-1} \\ g_{n-k} \end{array}$$

where $a = n - k + (n - k - 3) - (j - 1)$ ($j = n - k - 1, n - k - 2, \dots, 4, 3$). Defining an equivariant diffeomorphism $\varphi: (\mathbb{Z}_2, M(B_a)) \rightarrow (\mathbb{Z}_2, M(B_b))$ by $\varphi([z_2, \dots, z_n]) = [z_2, \dots, z_{j-1}, \mathbf{i}z_j, z_{j+1}, \dots, z_n]$, we obtain that (4.72) is equivalent to

$$A_{\ell b} = \left(\begin{array}{c|cccccccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4(b)} & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \{\hat{1}\}_{n(b)} \\ \hline & 1 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ & & 1 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ & & & \ddots & & & & & & & & \vdots \\ & & & & 1 & 1 & 0 & \dots & 0 & 0 & \dots & 0 \\ & & & & & 1 & 1 & \dots & 1 & 0 & \dots & 0 \\ & & & & & & 1 & \dots & 1 & 0 & \dots & 0 \\ & & & & & & \ddots & & & & & \vdots \\ & & & & & & & 1 & 1 & 0 & \dots & 0 \\ & & & & & & & & 1 & 1 & \dots & 1 \\ & & & & & & & & & & & I_k \end{array} \right) \begin{array}{l} g_1 \\ g_2 \\ g_3 \\ \vdots \\ g_{j-2} \\ g_{j-1} \\ g_j \\ \vdots \\ g_{n-k-1} \\ g_{n-k} \end{array}$$

where $b = n - k + (n - k - 3) - (j - 2)$. Therefore the previous Bott matrix is equivalent to

$$A_{\ell c} = \left(\begin{array}{c|cccccccc} 1 & 1 & 0 & \{\hat{1}\}_{4(c)} & \dots & \{\hat{1}\}_{n-k(c)} & \{\hat{1}\}_{n-k+1(c)} & \dots & \{\hat{1}\}_{n(c)} \\ \hline 0 & & & & B_{2(n-k)-4} & & & & \end{array} \right)$$

where $c = n - k + (n - k - 3) - 1$.

Finally each Bott matrix

$$\left(\begin{array}{c|cccc} 1 & 1 & 1 & \{\hat{1}\}_{4(i)} & \dots & \{\hat{1}\}_{n(i)} \\ \hline 0 & & & & B_j & \end{array} \right)$$

for $j = 2, \dots, 2(n - k) - 4$ is equivalent to one of Bott matrices in (4.22) or (4.36) by the equivariant diffeomorphism $\varphi([z_2, \dots, z_n]) = [\mathbf{i}z_2, z_3, \dots, z_n]$. This completes the proof of the theorem. $\square$

Acknowledgment. I would like to thank Professor Mikiya Masuda for his useful suggestions.

REFERENCES

- [1] M. Grossberg and Y. Karshon, *Bott towers, complete integrability, and the extended character of representations*, Duke Math. J 76 (1994) 23-58.
- [2] Y. Kamishima, and A. Nazra, *Seifert fibred structure and rigidity on real Bott towers*, 2009. To appear.
- [3] A. Nazra, *Real Bott Tower*, Tokyo Metropolitan University, Master Thesis 2008.
- [4] A. Nazra, *Geometry of Transformation Groups and Related Topics*, Diffeomorphism Type of Real Bott Towers, RIMS Kokyuroku 1612, Kyoto University, September 2008, 165-176.
- [5] Wolf, J.A., *Spaces of Constant Curvature*, McGraw-Hill Book Company, 1967.

DEPARTMENT OF MATHEMATICS, ANDALAS UNIVERSITY, KAMPUS UNAND LIMAU MANIS PADANG 25163, INDONESIA

E-mail address: `admi@fmipa.unand.ac.id`

Current address: Department of Mathematics, Tokyo Metropolitan University, Minami-Ohsawa 1-1, Hachioji, Tokyo 192-0397, JAPAN

E-mail address: `nazra-admi@ed.tmu.ac.jp`